

KUMMER COVERS AND BRAID MONODROMY

ENRIQUE ARTAL BARTOLO, JOSÉ IGNACIO COGOLLUDO-AGUSTÍN,
AND JORGE ORTIGAS-GALINDO

ABSTRACT. In this work we describe a method to reconstruct the braid monodromy of the preimage of a curve by a Kummer cover. This method is interesting, since it combines two techniques, namely, the reconstruction of a highly non-generic braid monodromy with a systematic method to go from a non-generic to a generic braid monodromy. This “*generification*” method is independent from Kummer covers and can be applied in more general circumstances since non-generic braid monodromies appear more naturally and are oftentimes much easier to compute. Explicit examples are computed using these techniques.

INTRODUCTION

A Kummer cover is a map $\pi_n : \mathbb{P}^2 \rightarrow \mathbb{P}^2$ given by $\pi_n([x : y : z]) := [x^n : y^n : z^n]$. Kummer covers are a very useful tool in order to construct complicated algebraic curves starting from simple ones. Since Kummer covers are finite Galois covers of $\mathbb{P}^2 \setminus \{xyz = 0\}$ with $\text{Gal}(\pi_n) \cong \mathbb{Z}/n\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$, topological properties of the new curves can be obtained: Alexander polynomial, fundamental group, characteristic varieties and so on (see [3, 4, 16, 34, 23, 18] for papers using these techniques).

On the other hand, the braid monodromy of a plane projective curve is a powerful invariant that provides a way to compute the fundamental group of its complement and was originally described as a formalization of the Zariski-van Kampen method [35, 24] (see [28, 21] also [17] and references therein for a detailed exposition on the subject). However, braid monodromy is a much more powerful invariant, since in fact it encodes the topology of the embedding of the curve, as well as the isomorphism problem for surfaces whose branching locus over \mathbb{P}^2 is a given curve (see [12, 26, 30, 14, 13, 25, 9, 20, 1] to name only a few).

In this work we focus our attention on a method to reconstruct the braid monodromy of the preimage of a curve by a Kummer cover. This method is interesting in and of itself, since it combines two techniques, namely, the reconstruction of a highly non-generic braid monodromy with a systematic method to go from a non-generic to a generic braid monodromy. This “*generification*” method is independent from Kummer covers and can be applied in more general circumstances (see also [31, 6, 2]). The reason for this is that oftentimes, non-generic braid monodromies are much easier to compute.

2010 *Mathematics Subject Classification.* 14F35, 14H50, 14F45, 57M12, 14E20.

Key words and phrases. Kummer covers, braid monodromy, plane curves.

All authors are partially supported by the Spanish Ministry of Education MTM2010-21740-C02-02.

The importance of finding theoretical arguments to obtain braid monodromies lies in the fact that any direct computation requires polynomial root approximation, whose exactness can only be justified in particularly simple cases, such as curves whose equation can be given in $\mathbb{Z}[x, y, z]$ ([11, 12]), strongly real curves ([7, p. 17] and references therein), or line arrangements (see [8, 32, 19, 33, 20]).

The layout of the paper is as follows: after describing in §1 the main objects to be used such as generic and non-generic braid monodromies, Kummer covers and extended braid monodromies are described in §2 as the main tool to recover a non-generic braid monodromy of a Kummer cover from a braid monodromy of the base. Useful *generification* techniques are described in §4. Sections §3 and §5 are more technical and describe useful systems of Artin generators for the braids obtained after a Kummer cover and the properties and singularities of Kummer transforms. Finally, a detailed account of numerous examples (some classical and some new) of the power of Kummer transforms is given in §6, where generic braid monodromies of smooth curves, sextics with six cusps (on a conic and otherwise), sextics with nine cusps, Hesse, Ceva, and MacLane arrangements are provided.

1. SETTINGS: BRAID MONODROMY

After the work of Zariski, braid monodromy was defined by Chisini [15] but it was necessary to wait for Moishezon [30] in order to get further applications of this invariant.

1.1. Construction.

Let us fix a curve $\bar{C} \subset \mathbb{P}^2$ of degree d , a point $P_y \in \mathbb{P}^2$ and a line \bar{L}_∞ such that $P_y \in \bar{L}_\infty$. We say that the curve is *horizontal* with respect to P_y if it does not contain any line through P_y ; we assume \bar{C} to be horizontal. We consider a system of coordinates $[X : Y : Z]$ such that $P_y := [0 : 1 : 0]$ and $\bar{L}_\infty := \{Z = 0\}$. We identify $\mathbb{C}^2 \equiv \mathbb{P}^2 \setminus \bar{L}_\infty$ with affine coordinates $(x, y) \equiv [x : y : 1]$.

Let $F(x, y, z) = 0$ be a reduced equation of \bar{C} , $k := \deg_y F$

$$F(x, y, z) = \sum_{j=0}^k \bar{a}_{d-j}(x, z)y^j, \quad \bar{a}_{d-k}(x, z) \neq 0, \quad \bar{a}_j \text{ homogeneous of degree } j,$$

normalized such that the coefficient of the term of higher degree of $\bar{a}_{d-k}(x, z)$ in x is 1. The fact that \bar{C} is horizontal is equivalent to $\gcd(F, \bar{a}_{d-k}) = 1$.

The pencil of lines through P_y is identified with $\mathbb{P}^1 \equiv \mathbb{C} \cup \{\infty\}$, where ∞ corresponds with \bar{L}_∞ . Following the previous notation the lines in the pencil are denoted by $\bar{L}_t := \{X - tZ = 0\}$. Let us restrict our attention to the affine part. Let $\mathcal{C} := \bar{C} \cap \mathbb{C}^2$ and $L_t := \bar{L}_t \cap \mathbb{C}^2$; the line L_t has equation $x = t$ while \mathcal{C} has equation $f(x, y) = 0$, where

$$f(x, y) := F(x, y, 1) = \sum_{j=0}^k a_{d-j}(x)y^j, \quad a_j(x) := \bar{a}_j(x, 1).$$

Let $\mathcal{B} := \{t \in \mathbb{C} \mid \#(L_t \cap \mathcal{C}) < k\}$; this is a finite set which contains the roots of $a_{d-k}(x)$ (if any) and the values t such that $L_t \not\subset \mathcal{C}$. The set \mathcal{B} is the zero locus of the product of $a_{d-k}(x)$ and the discriminant of $f(x, y)$ with respect to y .

Let $\Sigma_k(\mathbb{C}) := \{A \subset \mathbb{C} \mid \#A = k\}$ be a configuration space of \mathbb{C} ; for any $A := \{x_1, \dots, x_k\} \in \Sigma_k(\mathbb{C})$ the fundamental group $\pi_1(\Sigma_k(\mathbb{C}); A) =: \mathbb{B}(x_1, \dots, x_k)$ is isomorphic to the braid group \mathbb{B}_k with the usual Artin presentation

$$(1.1) \quad \mathbb{B}_k := \left\langle \sigma_1, \dots, \sigma_{k-1} \mid \begin{array}{l} [\sigma_i, \sigma_j] = 1, \\ 1 < i+1 < j < k \end{array} \quad \begin{array}{l} \sigma_i \cdot \sigma_{i+1} \cdot \sigma_i = \sigma_{i+1} \cdot \sigma_i \cdot \sigma_{i+1} \\ 1 \leq i < k-1 \end{array} \right\rangle.$$

For the next sections we need to describe a canonical identification between \mathbb{B}_k and $\mathbb{B}(x_1, \dots, x_k)$; the group $\pi_1(\Sigma_k(\mathbb{C}); A)$ is identified with the homotopy classes of sets of arcs $\varphi_1, \dots, \varphi_k : [0, 1] \rightarrow \mathbb{C}$ starting and ending in A and such that $\#\{\varphi_1(t), \dots, \varphi_k(t)\} = k, \forall t \in [0, 1]$. Let us order the points of A , say x_1, \dots, x_k and consider a set I of simple segments $A_i, 1 \leq i < k$, such that $\partial A_i = \{x_i, x_{i+1}\}$, $A_i \cap A_{i+1} = \{x_{i+1}\}$ and the other intersections are empty; such a collection I will be called a *diagram system* for (x_1, \dots, x_k) . Then we associate to σ_i the braid which is constant for $x_1, \dots, x_{i-1}, x_{i+2}, \dots, x_k$ and behaves in a small small closed (topological) disk $N(A_i)$ such that A_i is a diameter as follows. The points x_i and x_{i+1} counterclockwise exchange along $\partial N(A_i)$. These generators may be understood as *half-twists* around A_i .




FIGURE 1. Diagram system, $k = 5$.

There is also a basis of the free group $\pi_1(\mathbb{C} \setminus A; x_0)$ if one chooses a simple edge A_0 from x_0 to x_1 intersecting $\bigcup_{i=1}^{k-1} A_i$ only at x_1 . This basis μ_1, \dots, μ_k is obtained as follows: take small disks Δ_i centered at x_i and assume that their intersection with $A_{i-1} \cup A_i$ are diameters with ends x_i^-, x_i^+ . Then μ_i is defined as follows: take a path α_i from x_0 to x_i^- running along $A_0 \cup \dots \cup A_{i-1}$ outside the interior of the disk Δ_j and goes counterclockwise along $\partial \Delta_j$ from x_j^- to x_j^+ , $1 \leq j \leq i$. Let β_i be the closed path obtained by running counterclockwise along $\partial \Delta_i$ with base point x_i^- and define $\mu_i := \alpha_i \cdot \beta_i \cdot \alpha_i^{-1}$.

The group \mathbb{B}_k acts geometrically on the group $\mathbb{F}_k := \pi_1(\mathbb{C} \setminus A; x_0)$ as follows:

$$(1.2) \quad \mu_i^{\sigma_j} := \begin{cases} \mu_{i+1} & \text{if } j = i, \\ \mu_i \cdot \mu_{i-1} \cdot \mu_i^{-1} & \text{if } j = i-1, \\ \mu_i & \text{otherwise.} \end{cases}$$

Notation 1.1. As usual, $a^b = b^{-1}ab$ is the conjugation of a by b . Also, for short, we will write $b * a = bab^{-1}$.




FIGURE 2. Geometric version of the action of \mathbb{B}_k on \mathbb{F}_k .

There are two important facts in these definitions; the element $\mu_\infty := (\mu_k \cdots \mu_1)^{-1}$ is a meridian of the point at infinity and μ_∞ is a fixed point by the action of \mathbb{B}_k . We say that (μ_1, \dots, μ_k) is an *ordered geometric basis* of $\pi_1(\mathbb{C} \setminus A; x_0)$. As a general notation, if G is a group and $\mathbf{x} := (x_1, \dots, x_k) \in G^k$ we define the *pseudo-Coxeter element* of \mathbf{x} as $c_{\mathbf{x}} := x_k \cdots x_1$.

After this digression, note that f defines a map $\tilde{f} : \mathbb{C} \setminus \mathcal{B} \rightarrow \Sigma_k(\mathbb{C})$.

Definition 1.2. The *braid monodromy* of the triple $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ is the morphism

$$\nabla : \pi_1(\mathbb{C} \setminus \mathcal{B}; t_0) \rightarrow \mathbb{B}_k, \quad t_0 \in \mathbb{C} \setminus \mathcal{B},$$

defined by \tilde{f} on the fundamental group.

Remark 1.3. Consider a geometric basis $(\gamma_1, \dots, \gamma_r)$ of $\pi_1(\mathbb{C} \setminus \mathcal{B}; t_0)$ and let c_∞ be its pseudo-Coxeter element. Note that ∇ is determined by $(\nabla(\gamma_1), \dots, \nabla(\gamma_r)) \in \mathbb{B}_k^r$ having as pseudo-Coxeter element $\nabla(c_\infty)$.

The braid monodromy measures the motions of the points of \mathcal{C} along the affine lines L_t (identified with \mathbb{C}).

There are a lot of choices in order to obtain an element of \mathbb{B}_k^r from $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$. It is not hard to check that these choices are given by the orbits of an action of $\mathbb{B}_k \times \mathbb{B}_r$ on \mathbb{B}_k^r . The action of \mathbb{B}_k is given by simultaneous conjugation. The action of \mathbb{B}_r is defined as follows; let h_1, \dots, h_{r-1} an Artin system of generators of \mathbb{B}_r . Then, if $(\tau_1, \dots, \tau_r) \in \mathbb{B}_k^r$, then:

$$(1.3) \quad (\tau_1, \dots, \tau_r)^{h_i} := (\tau_1, \dots, \tau_{i-1}, \tau_{i+1}, \tau_{i+1} \cdot \tau_i \cdot \tau_{i+1}^{-1}, \tau_{i+2}, \dots, \tau_r);$$

h_i is called a *Hurwitz move*. In particular for a choice of $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ two objects are unique and well-defined: the conjugacy classes of the pseudo-Coxeter element and of the *monodromy group*, i.e., the group generated by $\nabla(\gamma_1), \dots, \nabla(\gamma_r)$.

In light of the previous discussion, a braid monodromy ∇ of a triple $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ will sometimes be considered as a morphism (see Definition 1.2) or as a list of braids $(\nabla(\gamma_1), \dots, \nabla(\gamma_r))$, where $\gamma_1, \dots, \gamma_r$ is a geometric basis.

1.2. Applications.

The first application of braid monodromy is the computation of the fundamental group, see [5].

Given ∇ the braid monodromy of a triple $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$. Consider $\mathbb{C}^2 := \mathbb{P}^2 \setminus \bar{L}_\infty$, $\mathcal{C} := \bar{\mathcal{C}} \cap \mathbb{C}^2$, and L_t ($t \in \mathcal{B}$) the non-generic fibers of the pencil of lines through P_y as above. One has the following result.

Theorem 1.4. *The group $\pi_1(\mathbb{C}^2 \setminus (\mathcal{C} \cup \bigcup_{t \in \mathcal{B}} L_t))$ has the following presentation:*

$$\left\langle \mu_1, \dots, \mu_k, \tilde{\gamma}_1, \dots, \tilde{\gamma}_r \left| \begin{array}{l} \tilde{\gamma}_i^{-1} \cdot \mu_j \cdot \tilde{\gamma}_i = \mu_j^{\nabla(\gamma_i)} \\ 1 \leq i \leq r; 1 \leq j \leq k \end{array} \right. \right\rangle.$$

A triple $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ (or simply a curve \mathcal{C}) is said to be *fully horizontal* if \mathcal{C} has no vertical asymptotes, i.e., if $b_{d-k}(x) = 1$. In that case, if $\mathcal{B}_0 \subset \mathcal{B}$ (maybe empty), corresponding to the generators $\{\gamma_{i_1}, \dots, \gamma_{i_s}\}$, one has a new version of Zariski-van Kampen Theorem, due to the fact that the above generators $\tilde{\gamma}_j$ are meridians of the corresponding lines.

Theorem 1.5. *The group $\pi_1(\mathbb{C}^2 \setminus (\mathcal{C} \cup \bigcup_{t \in \mathcal{B}_0} L_t))$ has the following presentation:*

$$\left\langle \mu_1, \dots, \mu_k, \tilde{\gamma}_{i_1}, \dots, \tilde{\gamma}_{i_s} \left| \begin{array}{ll} \tilde{\gamma}_{i_j}^{-1} \cdot \mu_\ell \cdot \tilde{\gamma}_{i_j} = \mu_\ell^{\nabla(\gamma_{i_j})}, & \mu_j = \mu_j^{\nabla(\gamma_i)} \\ 1 \leq j \leq s; 1 \leq \ell \leq k & i \neq i_1, \dots, i_s; 1 \leq j \leq k \end{array} \right. \right\rangle.$$

There is also a statement like Theorem 1.5 for curves with vertical asymptotes but it is more technical, see [12] or [28].

Let us assume now that $P_y \notin \bar{\mathcal{C}}$. In this case the triple $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ is automatically fully horizontal and $d = k$. In this case another version can be stated, which is closer to the original Zariski-van Kampen Theorem.

Theorem 1.6. *The group $\pi_1(\mathbb{P}^2 \setminus \bar{\mathcal{C}})$ has the following presentation:*

$$\left\langle \mu_1, \dots, \mu_d \left| \begin{array}{l} \mu_j = \mu_j^{\nabla(\gamma_i)}, \\ 1 \leq i \leq r; 1 \leq j \leq d \end{array} \right. \mu_d \cdot \dots \cdot \mu_1 = 1 \right\rangle.$$

There is also a procedure to compute $\pi_1(\mathbb{P}^2 \setminus \bar{\mathcal{C}})$ when $P_y \in \bar{\mathcal{C}}$, but the general formula is not so closed since it depends heavily on the local singularity $(\bar{\mathcal{C}} \cup \bar{L}_\infty, P_y)$.

1.3. Generic braid monodromies.

We finish this sequence of versions of Zariski-van Kampen Theorem with the generic case. We assume $P_y \notin \bar{\mathcal{C}}$, $\bar{L}_\infty \cap \bar{\mathcal{C}}$ (i.e they intersect at d distinct points), and moreover for each $t \in \mathcal{B}$ there is exactly one point $P_t \in \bar{L}_t \cap \bar{\mathcal{C}}$ where the intersection is not transversal and satisfies

$$(1.4) \quad (\bar{\mathcal{C}} \cdot \bar{L}_t)_{P_t} = \begin{cases} 2 & \text{if } (\bar{\mathcal{C}}, P_t) \text{ is smooth,} \\ m_t & \text{if } (\bar{\mathcal{C}}, P_t) \text{ is singular,} \end{cases}$$

where m_t is the multiplicity of the germ $(\bar{\mathcal{C}}, P_t)$. In the singular case, it means that \bar{L}_t is not in the tangent cone of $(\bar{\mathcal{C}}, P_t)$ and, in the smooth case, that P_t is not an inflection point. In order to state the final form of Zariski-van Kampen Theorem, we need some notation. The above conditions imply that for each γ_i , $1 \leq i \leq r$, we can express $\nabla(\gamma_i) = \eta_i \cdot \tau_i \cdot \eta_i^{-1}$, where $\eta_i, \tau_i \in \mathbb{B}_d$ and τ_i is a positive word in $\Sigma_i := \{\sigma_j\}_{j \in s_i}$, where $s_i \subset \{1, \dots, d-1\}$ and $\#s_i = (\bar{\mathcal{C}} \cdot \bar{L}_t)_{P_t} - 1$.

For each i , $(1 \leq i \leq r)$ the elements $\mu_j(i) := \mu_j^{(\eta_i^{-1})}$ represent another basis of \mathbb{F}_d . Let us denote by $\mu_j(i)^{\sigma_k(i)}$ the action (1.2) written in the basis $\mu_1(i), \dots, \mu_d(i)$.

Theorem 1.7. *The group $\pi_1(\mathbb{P}^2 \setminus \bar{\mathcal{C}})$ has the following presentation:*

$$\left\langle \mu_1, \dots, \mu_d \left| \begin{array}{l} \mu_j(i) = \mu_j(i)^{\tau_i(i)}, \quad \mu_d \cdot \dots \cdot \mu_1 = 1 \\ 1 \leq i < r; \quad j \in s_i \end{array} \right. \right\rangle.$$

In this case, the pseudo-Coxeter element of $(\nabla(\gamma_1), \dots, \nabla(\gamma_r))$ is Δ_d^2 , the positive generator of the central element of \mathbb{B}_d . This is why in the literature braid monodromy is also referred to as a *factorization* of Δ_d^2 . The main point is that under these genericity conditions, braid monodromy is an invariant of $\bar{\mathcal{C}}$ independent of the (generic) choice of P_y and \bar{L}_∞ . Moreover, if we can connect two curves by a path of equisingular curves, then they share the braid monodromy.

A more general type of genericity occurs simply when $P_y \notin \bar{\mathcal{C}}$, $\bar{L}_\infty \pitchfork \bar{\mathcal{C}}$. We will refer to such a triple as *generic at infinity*. In this case Theorem 1.7 is still true, but the sets s_i have to be replaced by a finite number of disjoint subsets $s_{i,1}, \dots, s_{i,\kappa_i} \subset \{1, \dots, d-1\}$.

An interesting case of genericity at infinity occurs when condition (1.4) holds, but each fiber L_t , $t \in \mathcal{B}$ is allowed to have more than one ramification point. We refer to this case as *local genericity*. From a locally generic braid monodromy, a generic braid monodromy might easily be reconstructed (see Proposition 4.3).

2. PENCILS OF LINES AND KUMMER COVERS

Following notation from §1, let us fix a curve $\bar{\mathcal{C}} \subset \mathbb{P}^2$ of degree k . We assume that $P_y := [0 : 1 : 0] \notin \bar{\mathcal{C}}$. Let $\bar{\mathcal{C}}_n := \pi_n^{-1}(\bar{\mathcal{C}})$ be its Kummer transform. Denote by $\bar{L}_t := \{X - tZ = 0\}$ the pencil of lines through P_y .

Remark 2.1. Most of the results of this section also hold if $P_y \in \bar{\mathcal{C}}$ and the tangent cone of $(\bar{\mathcal{C}}, P_y)$ is $Z = 0$ (or alternatively, if a change of variables can be performed to match these conditions).

Let $\bar{L}_Y := \{Y = 0\} \subset \mathbb{P}^2$ and let L_Y be its affine part. We consider the set of lines through P_y which are not transversal to $\bar{\mathcal{C}} \cup \bar{L}_Y$ and to which we include the lines \bar{L}_0 and \bar{L}_∞ if necessary (due to their special relation with π_n). Let

$$B := \{t \in \mathbb{P}^1 \mid \bar{L}_t \not\pitchfork (\bar{\mathcal{C}} \cup \bar{L}_Y)\}, \quad B^* := B \cap \mathbb{C}^*, \quad \tilde{B} := B^* \cup \{0\}, \quad B^\infty := B \cup \{\infty\}.$$

The braid monodromy of $\bar{\mathcal{C}} \cup \bar{L}_Y$ with respect to P_y and \bar{L}_∞ is the morphism

$$\nabla : \pi_1(\mathbb{C} \setminus B; t_0) \rightarrow \mathbb{B}_{k+1}, \quad t_0 \in \mathbb{C} \setminus \tilde{B}.$$

Note that the image of ∇ is contained in $\mathbb{B}_{k,1}$, the subgroup of \mathbb{B}_{k+1} given by all the braids whose associated permutation fixes a given point (see §3 for a description of this group).

Definition 2.2. The map

$$(2.1) \quad \tilde{\nabla} : \pi_1(\mathbb{C} \setminus \tilde{B}; t_0) \rightarrow \mathbb{B}_{k,1}$$

is called the *extended braid monodromy* of the triple $(\bar{\mathcal{C}}, P_y, L_\infty)$ with respect to L_Y .

Note that if $0 \notin B$ then the image by $\tilde{\nabla}$ of a meridian around 0 is the trivial braid.

We define the analogous objects for $\bar{\mathcal{C}}_n$: $B_n, B_n^*, \tilde{B}_n, B_n^\infty, \nabla_n, \tilde{\nabla}_n$ for $t_{0,n}$, an n -th root of t_0 . The following Lemmas are easy consequences of the properties of a cover.

Lemma 2.3. *Let $\phi_n : \mathbb{C} \rightarrow \mathbb{C}$ be the map defined by $\phi_n(t) := t^n$. Then, $B_n^* = \phi_n^{-1}(B^*)$ and $\tilde{B}_n = \phi_n^{-1}(\tilde{B})$.*

Lemma 2.4. *The map π_n induces degree n coverings $\varphi_n : L_t \mapsto L_{t^n}$. Moreover, the preimage of a line L_s by π_n is the disjoint union of L_t , $t^n = s$, each one inducing a covering φ_n as above.*

As a consequence of these lemmas, the braid monodromy $\tilde{\nabla}_n$ can be thought as follows: consider the loops γ in $\mathbb{C} \setminus \tilde{B}$ which can be lifted as loops by ϕ_n and replace the braid $\nabla(\gamma)$, where k points move in \mathbb{C}^* , by the braid obtained by the constant string 0 and the n -roots of these k points. This construction defines a map

$$\tilde{\rho}_{n,k} : \mathbb{B}_{k,1} \rightarrow \mathbb{B}_{nk,1}$$

which is described in §3.

Let us summarize the results. If $r := \#B^*$, we denote by \mathbb{F}_{r+1} the free group $\pi_1(\mathbb{C} \setminus \tilde{B}; t_0)$. Analogously, we denote by \mathbb{F}_{nr+1} the free group $\pi_1(\mathbb{C} \setminus \tilde{B}_n; t_{0,n})$. Using covering theory, the map $(\phi_n)_*$ fits in a short exact sequence

$$0 \rightarrow \mathbb{F}_{nr+1} \rightarrow \mathbb{F}_{r+1} \rightarrow \mathbb{Z}/n\mathbb{Z} \rightarrow 0.$$

The following diagram holds:

$$(2.2) \quad \begin{array}{ccc} \mathbb{F}_{r+1} & \xrightarrow{\tilde{\nabla}} & \mathbb{B}_{k,1} \\ \uparrow & & \downarrow \\ \mathbb{F}_{nr+1} & \xrightarrow{\tilde{\nabla}_n} & \mathbb{B}_{nk,1} \end{array}$$

Using this diagram, one can recover geometric bases and relate the central elements of the corresponding braid groups. The following makes this construction useful.

Summarizing, let $\tilde{\nabla} := (\tau_1, \dots, \tau_{r+1}) \in \mathbb{B}_{k,1}^{r+1}$ be an extended braid monodromy for $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ with respect to L_Y , and τ_{r+1} is the braid corresponding to \bar{L}_0 . Consider $\pi_n(X : Y : Z) = [X^n : Y^n : Z^n]$ the n -th Kummer cover of \mathbb{P}^2 and denote by $\bar{\mathcal{C}}_n$ the preimage of $\bar{\mathcal{C}}$ by π_n . The braid monodromy $\tilde{\nabla}$ produces a list of braids $\tilde{\nabla}_n \in \mathbb{B}_{nk,1}^{nr+1}$ as described in diagram (2.2). One has the following

Proposition 2.5. *The element $\tilde{\nabla}_n$ described above is an extended braid monodromy for the triple $\pi_n^{-1}(\bar{\mathcal{C}}, P_y, \bar{L}_\infty) = (\bar{\mathcal{C}}_n, P_y, \bar{L}_\infty)$ with respect to $\pi_n^{-1}(\bar{L}_Y) = \bar{L}_Y$.*

Proof. It is an immediate consequence of the previous construction and Lemmas 2.3 and 2.4. \square

Remark 2.6. Note that the braid monodromy obtained from $\tilde{\nabla}_n$ via the forgetful map $\mathbb{B}_{nk,1} \rightarrow \mathbb{B}_{nk}$ is (for the interesting cases) a highly non-generic braid monodromy.

Lemma 2.7. *Let $(\gamma_1, \dots, \gamma_{r+1})$ be a geometric basis of \mathbb{F}_{r+1} such that γ_{r+1} is a meridian of $0 \in \tilde{B}$. Then, the ordered list*

$$(v_0, \dots, v_j, \dots, v_{n-1}, \gamma_{r+1}^n),$$

where $v_j := (\gamma_1^{\gamma_j^{r+1}}, \dots, \gamma_r^{\gamma_j^{r+1}})$ (see Notation 1.1), forms a geometric basis of \mathbb{F}_{nr+1} .

Lemma 2.8. *Let $\Delta_{k+1}^2 \in \mathbb{B}_{k,1}$ be the positive generator of the center of \mathbb{B}_{k+1} . Then $\Delta_{k+1}^{2n} = \Delta_{nk+1}^2$ via the inclusion $\mathbb{B}_{k,1} \hookrightarrow \mathbb{B}_{nk,1}$.*

The purpose of the upcoming sections will be to describe an effective way to obtain a generic braid monodromy for $\tilde{\mathcal{C}}_n$ from the given computation of $\tilde{\nabla}_n$. This is an outline of the general strategy, which will be developed in what follows:

- If $\bar{L}_Y \not\subseteq \bar{\mathcal{C}}_n$, we compose $\tilde{\nabla}_n$ with the natural map $\mathbb{B}_{nk,1} \rightarrow \mathbb{B}_{nk}$ obtained by forgetting the constant string 0.
- If $0 \notin B_n$ then the map factorizes through $\mathbb{F}_{nr+1} \rightarrow \mathbb{F}_{nr}$ whose kernel is generated by the meridians around 0.
- If $\infty \in B_n$ we change the line at infinity; the image α of a meridian at infinity is obtained as follows. Let τ be the pseudo-Coxeter of the braid monodromy; then $\alpha := \Delta^2 \tau^{-1}$, where Δ^2 is the positive generator of the center of the braid group.
- Slightly move the projection point P_y to obtain a generic projection.

3. THE MOTION OF n -ROOTS

As was introduced in §2, let $\mathbb{B}_{k,1}$ be the subgroup of \mathbb{B}_{k+1} given by all the braids whose associated permutation fixes a given point in \mathbb{C} , say 0.

We identify \mathbb{B}_{k+1} with the group of braids with ends in $k+1$ non-negative real points $x_1 > \dots > x_k > x_{k+1} = 0$ and we consider the Artin generators $\sigma_1, \dots, \sigma_k$ in \mathbb{B}_{k+1} which are geometrically associated with the diagram system obtained from the paths joining these points in the real line, see §1. Recall that any choice of a *diagram system* I of $k+1$ points in \mathbb{C} induces an Artin system of generators (1.1) of \mathbb{B}_{k+1} . Note that the only half-twist σ_i that moves 0 is σ_k . The following result is well known.

Lemma 3.1. *The group $\mathbb{B}_{k,1}$ is generated by $\sigma_1, \dots, \sigma_{k-1}, \sigma_k^2$ (as a subgroup of \mathbb{B}_{k+1}) and it is naturally isomorphic to the group of braids of k strands in \mathbb{C}^* .*

Let us consider the map $\rho_n : \mathbb{C}^* \rightarrow \mathbb{C}^*$, given by $\rho_n(z) := z^n$. Note that this map induces a morphism $\tilde{\rho}_n : \mathbb{B}_{k,1} \rightarrow \mathbb{B}_{nk,1}$ via the multivalued function ρ_n^{-1} . The goal of this section is to give explicit formulæ for this morphism. In order to do so, one needs to explicitly choose systems of generators for \mathbb{B}_{k+1} and \mathbb{B}_{nk+1} .

Consider the diagram system I for $\mathbb{B}_{k,1}$ described above. The image of the half twist σ_i , associated with A_i , $1 \leq i < k$, is a product of half-twists associated with the pairwise disjoint arcs $A_{i,1}, \dots, A_{i,n}$ such that $\rho_n^{-1}(A_i) = A_{i,1} \cup \dots \cup A_{i,n}$. The image of σ_k^2 is more complicated since $A_{k,1}, \dots, A_{k,n}$ intersect at 0. However, note that the natural system of arcs $A_{i,j}$ does not produce a diagram system for \mathbb{B}_{nk+1} . In order to solve this situation one needs to define diagram systems for $\mathbb{B}_{nk,1}$ that allow to describe the arcs $A_{i,j}$ naturally. For different purposes, different diagram systems might be more suited. Here we will concentrate our attention in two particular diagram systems: the circular and the radial.


3.1. Circular diagram systems.

Let us denote $d := nk$. We identify \mathbb{B}_{d+1} with $\mathbb{B}(\{\xi_n^j t_i \mid 1 \leq i \leq k, 1 \leq j \leq n\} \cup \{0\})$, where t_i is the non-negative n -th root of $x_i \in \mathbb{R}_{\geq 0}$ and $\xi_n := \exp(\frac{2\pi\sqrt{-1}}{n})$. Consider the following arcs:

- $c_{i,j}$ is the counterclockwise arc joining $t_i \xi_n^{j-1}$ to $t_i \xi_n^j$ in the circle centered at 0, $1 \leq j < n$;
- $c_{i,n}$ is the segment joining $t_i \xi_n^{n-1}$ and t_{i+1} .

These arcs and segments are illustrated in Figure 3(a). The list of arcs $\{c_{i,j}\}_{i,j}$ with a left-lexicographic order produces a diagram system called the *circular diagram system*. The half twist produced by an arc $c_{i,j}$ will be denoted by $\sigma_{i,j} \in \mathbb{B}_{nk,1}$. The half-twists associated with $A_{i,j}$ will be denoted by $\alpha_{i,j}$.

Figure 3(b) is obtained from Figure 3(a) after unwinding the circular diagram system into a straight line to get the usual projections. This will help visualize the rewriting of the half-twists $\alpha_{i,j}$ in terms of the $\sigma_{i,j}$.



(a) Circular diagram system of \mathbb{B}_{nk+1} for $k = n = 3$

(b) A *straightened* view of Figure 3(a).

FIGURE 3.

Consider $(i_1, j_1) < (i_2, j_2)$, where $<_\ell$ represents the left-lexicographic order (that is, either $i_1 < i_2$ or $i_1 = i_2$ and $j_1 < j_2$). We will define

$$(3.1) \quad \begin{aligned} P_{i_1, j_1}^{i_2, j_2} &:= \sigma_{i_1, j_1} \cdot \dots \cdot \sigma_{i_2, j_2} = \prod_{(i_1, j_1) \leq_\ell v \leq_\ell (i_2, j_2)} \sigma_v \\ \bar{P}_{i_1, j_1}^{i_2, j_2} &:= \sigma_{i_1, j_1}^{-1} \cdot \dots \cdot \sigma_{i_2, j_2}^{-1} = \prod_{(i_1, j_1) \leq_\ell v \leq_\ell (i_2, j_2)} \sigma_v^{-1} \\ M_{i_1, j_1}^{i_2, j_2} &:= \sigma_{i_2, j_2} \cdot \dots \cdot \sigma_{i_1, j_1} = \prod_{(i_2, j_2) \geq_\ell v \geq_\ell (i_1, j_1)} \sigma_v \\ \bar{M}_{i_1, j_1}^{i_2, j_2} &:= \sigma_{i_2, j_2}^{-1} \cdot \dots \cdot \sigma_{i_1, j_1}^{-1} = \prod_{(i_2, j_2) \geq_\ell v \geq_\ell (i_1, j_1)} \sigma_v^{-1}. \end{aligned}$$

Remark 3.2. Note that $P_{i_1, j_1}^{i_2, j_2} = (\bar{M}_{i_1, j_1}^{i_2, j_2})^{-1}$, $(\bar{P}_{i_1, j_1}^{i_2, j_2})^{-1} = M_{i_1, j_1}^{i_2, j_2}$. Also, note that

$$P_{i_1, j_1}^{i_2, j_2} * \sigma_{i_2, j_2} = \bar{M}_{i_1, j_1}^{i_2, j_2} * \sigma_{i_1, j_1} \quad (\text{resp. } \bar{P}_{i_1, j_1}^{i_2, j_2} * \sigma_{i_2, j_2}^{-1} = M_{i_1, j_1}^{i_2, j_2} * \sigma_{i_1, j_1}^{-1})$$

represents a half-twist (resp. a negative half-twist) interchanging $t_{i_1} \xi_n^{j_1-1}$ and $t_{i_2} \xi_n^{j_2+1}$ along a spiral arc. The arc corresponding to $P_{1,1}^{3,1} * \sigma_{3,1} = \bar{M}_{1,1}^{3,1} * \sigma_{1,1}$ (or $\bar{P}_{1,1}^{3,1} * \sigma_{3,1}^{-1} = M_{1,1}^{3,1} * \sigma_{1,1}^{-1}$) is shown in Figure 3(a).

Lemma 3.3. *Under the above conditions and for any $(i_1, j_1) < (i, j) < (i_2, j_2)$ one has*

- (1) $P_{i_1, j_1}^{i_2, j_2} * \sigma_{i_2, j_2} = \left(\bar{M}_{(i, j)^+}^{i_2, j_2} \cdot P_{i_1, j_1}^{i, j} \right) * \sigma_{i, j}$, where $(i, j)^+$ denotes the element following (i, j) in the left-lexicographic order.
- (2) $\bar{P}_{i_1, j_1}^{i_2, j_2} * \sigma_{i_2, j_2} = \left(M_{(i, j)^+}^{i_2, j_2} \cdot \bar{P}_{i_1, j_1}^{i, j} \right) * \sigma_{i, j}$.
- (3) $\alpha_{i, j} = \left(P_{i+1, j}^{k, n} \bar{P}_{i, j}^{k, n} \right) * \sigma_{k, n} = \left(P_{i+1, j}^{k, n} \sigma_{k, n}^2 M_{(i_2, j_2)^+}^{k, n-1} \bar{P}_{i, j}^{i_2, j_2} \right) * \sigma_{i_2, j_2} = \left(P_{i+1, j}^{k, n-1} \sigma_{k, n}^2 M_{i, j}^{k, n-1} \right) * \sigma_{i, j}$.

Proof. Property ((1)) is immediate by induction since

$$\begin{aligned} P_{i_1, j_1}^{i_2, j_2} * \sigma_{i_2, j_2} &= \left(P_{i_1, j_1}^{(i_2, j_2)-2} \cdot \sigma_{(i_2, j_2)^-} \right) * \sigma_{i_2, j_2} = \\ &= \left(P_{i_1, j_1}^{(i_2, j_2)-2} \cdot \sigma_{i_2, j_2}^{-1} \right) * \sigma_{(i_2, j_2)^-} = \left(\sigma_{i_2, j_2}^{-1} \cdot P_{i_1, j_1}^{(i_2, j_2)^-} \right) * \sigma_{(i_2, j_2)^-}, \end{aligned}$$

where $(i_2, j_2)^-$ denotes the element immediately smaller than (i_2, j_2) and $(i_2, j_2) - 2$ denotes its second predecessor $((i_2, j_2)^-)^-$. Property ((2)) is analogous.

In order to show Property ((3)) note that $\alpha_{i, j}$ is a half-twist that exchanges $t_{i_1} \xi_n^{j_1-1}$ and $t_{i+1} \xi_n^{j-1}$. Therefore, according to Remark 3.2 one has

$$\alpha_{i, j} = \left(P_{i+1, j}^{k, n} * \sigma_{k, n} \right) \cdot \left(\bar{P}_{i, j}^{k, n} * \sigma_{k, n} \right) \cdot \left(P_{i+1, j}^{k, n} * \sigma_{k, n}^{-1} \right).$$

A similar induction argument shows that

$$\left(P_{i+1, j}^{k, n} * \sigma_{k, n} \right) \cdot \left(\bar{P}_{i, j}^{k, n} * \sigma_{k, n} \right) \cdot \left(P_{i+1, j}^{k, n} * \sigma_{k, n}^{-1} \right) = \left(P_{i+1, j}^{k, n} \bar{P}_{i, j}^{k, n} \right) * \sigma_{k, n}.$$

The other equalities are a consequence of Remark 3.2 and Properties ((1)) and ((2)). \square

Proposition 3.4. *The map $\tilde{\rho}_n : \mathbb{B}_{k,1} \rightarrow \mathbb{B}_{d,1}$ is given by*

$$\begin{aligned} \sigma_i &\mapsto \prod_{j=1}^n \alpha_{i, j}, \quad i \in 1, \dots, k-1 \\ \sigma_k^2 &\mapsto \sigma_{k, n}^2 \sigma_{k, n-1} \sigma_{k, n-2} \dots \sigma_{k, 1} \end{aligned}$$

Proof. As mentioned at the beginning of this section, if $i \neq k$, the preimage of A_i by ρ_n is a disjoint union of segments $A_{i,1}, \dots, A_{i,n}$. Therefore the image of the half-twist σ_i associated with A_i is the product of the n half-twists $\alpha_{i,1} \cdot \dots \cdot \alpha_{i,n}$. Note that σ_k^2 corresponds to a full turn




FIGURE 4. Image of σ_k^2 , $n = 3$.

around 0, its preimage by ρ_n being a counterclockwise rotation of angle $\frac{2\pi}{n}$ of the points $\xi_n^j t_k$. This is nothing but $\sigma_{k,n}^2 \cdot \sigma_{k,n-1} \cdot \dots \cdot \sigma_{k,1}$ (see Figure 4). \square

Example 3.5. Let us consider the case $k = 1$ and the composition of $\tilde{\rho}_n$ with the natural projection of $\mathbb{B}_{n,1} \rightarrow \mathbb{B}_k$ given by forgetting the string 0. Then we have a map $\hat{\rho}_n : \mathbb{B}_{1,1} \rightarrow \mathbb{B}_n$ such that $\hat{\rho}_n(\sigma_1^2) = \sigma_{n-1} \sigma_{n-2} \cdot \dots \cdot \sigma_1$.

Example 3.6. Next consider the case $k = 2$ and the map $\hat{\rho}_n : \mathbb{B}_{2,1} \rightarrow \mathbb{B}_{2n}$ as in Example 3.5. Then, according to Lemma 3.3((3))

$$\hat{\rho}_n(\sigma_1) = \prod_{j=1}^n \left(P_{2,j}^{2,n-1} M_{1,j}^{2,n-1} \right) * \sigma_{1,j} = \prod_{j=1}^n (\sigma_{2,j} \cdot \dots \cdot \sigma_{2,n-1}^2 \cdot \dots \cdot \sigma_{1,j-1}) * \sigma_{1,j}$$

and

$$\hat{\rho}_n(\sigma_2^2) = \sigma_{2,n-1} \cdot \dots \cdot \sigma_{2,1}.$$

3.2. Radial diagram systems.

For some applications it is more useful to consider a different diagram system which will be called *radial*, see Figure 5(a).

We call the Artin generators for this system $\tilde{\sigma}_{i,j}$, $1 \leq i \leq k$, $1 \leq j \leq n$. According to this choice, these are the half-twists $\alpha_{i,j}$ associated with the segments $A_{i,j}$ for $1 \leq i \leq k-1$, that is,

$$(3.2) \quad \tilde{\sigma}_{i,j} = \alpha_{i,j}.$$

However, the element $\tilde{\sigma}_{k,j}$ corresponds to an arc joining $t_k \xi_n^j$ with $t_1 \xi_n^{j+1}$ as shown in Figure 5(a). Denote $\beta_j := \tilde{\sigma}_{k-1,j} \cdot \dots \cdot \tilde{\sigma}_{1,j}$. Then we have

$$(3.3) \quad \sigma_{k,j} = \beta_{j+1} \tilde{\sigma}_{k,j} \beta_{j+1}^{-1} \quad (1 \leq j \leq n-1)$$




FIGURE 5.

Moreover, $\sigma_{k,n} = \tilde{\sigma}_{k,n}$. The following result holds.

Proposition 3.7. *The map $\tilde{\rho}_n : \mathbb{B}_{k,1} \rightarrow \mathbb{B}_{d,1}$ is given by*

$$\begin{aligned} \sigma_i &\mapsto \prod_{j=1}^n \tilde{\sigma}_{i,j}, \quad 1 \leq i \leq k-1 \\ \sigma_k^2 &\mapsto \tilde{\sigma}_{k,n}^2 \prod_{(k-1,n) \geq_r v \geq_r (k,1)} \tilde{\sigma}_v \cdot \prod_{j=2}^n \beta_j^{-1}, \end{aligned}$$

where \geq_r represents the right-lexicographic order.


Proof. The result follows from Proposition 3.4. The formula for the image of σ_i is a direct consequence of (3.2). The formula for the image of σ_k^2 can be obtained using (3.3) and the fact that β_j commutes with $\beta_{j'}$ and with $\tilde{\sigma}_{i,j'}$ for $j' \leq j$ and $(i, j') \neq (k, j-1)$ (see Figure 6). \square

Example 3.8. Let us consider the case $n = 2$ and $\hat{\rho}_2 : \mathbb{B}_{k,1} \rightarrow \mathbb{B}_{2k}$ the composition of $\tilde{\rho}$ with the forgetful map as in Example 3.5. According to Proposition 3.7 one has

$$\begin{aligned} \sigma_i &\mapsto \tilde{\sigma}_{i,1} \tilde{\sigma}_{i,2}, \quad 1 \leq i \leq k-1 \\ \sigma_k^2 &\mapsto (\tilde{\sigma}_{k-1,2} \cdot \dots \cdot \tilde{\sigma}_{1,2}) * \tilde{\sigma}_{k,1} \end{aligned}$$

In fact, a more convenient description of these maps can be given for another choice of generators $\hat{\sigma}_i$ coming from the natural diagram system on the real line joining the preimages of x_k (which they all lie on the real line since $n = 2$).

$$\begin{array}{llll} \rho'_2 : \mathbb{B}_{k,1} & \rightarrow & \mathbb{B}_{2k+1} & \hat{\rho}'_2 : \mathbb{B}_{k,1} \rightarrow \mathbb{B}_{2k} \\ \sigma_i & \mapsto & \hat{\sigma}_i \hat{\sigma}_{2k-i+1} & \sigma_i \mapsto \hat{\sigma}_i \hat{\sigma}_{2k-i} \\ \sigma_k^2 & \mapsto & \hat{\sigma}_{k+1} \hat{\sigma}_k \hat{\sigma}_{k+1} & \sigma_k^2 \mapsto \hat{\sigma}_k. \end{array}$$

FIGURE 6. Image of σ_k^2 by Proposition 3.7.

4. DEFORMATIONS OF BRAID MONODROMIES

We are mostly interested in computing generic braid monodromies of curves (see §1.3). However, it is usually easier (and oftentimes more natural) to compute non-generic braid monodromies. For instance, those obtained after Kummer covers (even if the starting monodromy was generic) are typically non generic. In this section we show how to deal with non-generic braid monodromies.

4.1. From fully horizontal to generic at infinity.

Let us consider the braid monodromy of a fully horizontal triple $(\bar{\mathcal{C}}, P, \bar{L})$, where $P \notin \bar{\mathcal{C}}$ (e.g. the hypothesis in Theorem 1.6), where \bar{L} and $\bar{\mathcal{C}}$ are not necessarily transversal.

Proposition 4.1. *Assume $\bar{L} \not\lhd \bar{\mathcal{C}}$, let $(\tau_1, \dots, \tau_r) \in \mathbb{B}_d^r$ be a braid monodromy factorization for $(\bar{\mathcal{C}}, P, \bar{L})$, and consider a line \bar{L}' such that $P \in \bar{L}'$ and $\bar{L}' \lhd \bar{\mathcal{C}}$. Then $(\tau_1, \dots, \tau_r, \Delta_d^2(\tau_r \cdots \tau_1)^{-1})$ is a braid monodromy factorization for $(\bar{\mathcal{C}}, P, \bar{L}')$.*

Proof. Note that $\mathcal{B}' = \mathcal{B} \cup \{t_L\}$, where \mathcal{B} (resp. \mathcal{B}') is the ramification set for $(\bar{\mathcal{C}}, P, \bar{L})$ (resp. for $(\bar{\mathcal{C}}, P, \bar{L}')$). Since the line \bar{L}' is transversal, the pseudo-Coxeter of its braid monodromy factorization is Δ_d^2 and the result follows. \square

This will be a common situation when studying Kummer covers for curves which are not transversal to the axes. In that case we can be more explicit.

Let $T := (\tau_1, \dots, \tau_r, \tau_{r+1}) \in \mathbb{B}_{k,1}^{r+1}$ be an extended braid monodromy factorization for $(\bar{\mathcal{C}}, P_y = [0 : 1 : 0], \bar{L}_\infty)$ with respect to \bar{L}_Y . Assume that $\bar{L}_0 = \{X = 0\}$ is not transversal to $\bar{\mathcal{C}}$, and that τ_{r+1} is the braid corresponding to \bar{L}_0 . Let us denote by c_T its pseudo-Coxeter element.

Consider $\pi_n(X : Y : Z) = [X^n : Y^n : Z^n]$ the n -th Kummer cover of \mathbb{P}^2 and denote by $\bar{\mathcal{C}}_n$ the preimage of $\bar{\mathcal{C}}$ by π_n . The extended braid monodromy T produces an extended braid monodromy $T_n \in \mathbb{B}_{nk,1}^{nr+1}$ for $(\bar{\mathcal{C}}_n, \pi_n^{-1}P_y, \bar{L}_\infty)$ (see Proposition 2.5).

Proposition 4.2. *Under the above conditions, a generic at infinity extended braid monodromy factorization is obtained by adding to T_n the braid $\Delta_{k+1}^{2n} c_T^{-n}$, using the inclusion $\mathbb{B}_{k,1} \hookrightarrow \mathbb{B}_{nk,1}$ (see diagram (2.2)).*

Proof. It is a consequence of Proposition 4.1 and Lemma 2.8. \square

4.2. From locally generic to generic.

As announced in §1, some types of non-generic braid monodromies can be easily deformed into generic braid monodromies. This is the case with locally generic triples. Let $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ be a locally generic triple (see §1.3) and \mathcal{B} the set of ramification values of the projection. As already mentioned in §1.3, for each $t \in \mathcal{B}$, there exists a finite number of points $P_{t,1}, \dots, P_{t,\kappa_t} \in L_t$ satisfying (1.4).

Proposition 4.3. *Under the above conditions, let τ_t be the braid associated with L_t , $t \in \mathcal{B}$. Then there is a factorization $\tau_t = \tau_{t,\kappa_t} \cdot \dots \cdot \tau_{t,1}$, where $\tau_{t,1}, \dots, \tau_{t,\kappa_t}$ are pairwise commuting braids and each one corresponds to one non-transversal point in L_t .*

Moreover, replacing τ by $(\tau_{t,1}, \dots, \tau_{t,\kappa_t})$ (the order does not matter) in the braid monodromy factorization of $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ for each $t \in \mathcal{B}$ produces a generic braid monodromy factorization of $\bar{\mathcal{C}}$.

Proof. Since $P_y \notin \bar{\mathcal{C}}$ the triple $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ is fully horizontal, that is, there are no vertical asymptotes. Consider $P_{t,1}, \dots, P_{t,\kappa_t} \in L_t$. Due to the conic structure of $(\mathbb{C}^2, \mathcal{C})_{P_{t,i}}$ one can find disjoint Milnor polydisks $\mathbb{B}_{P_{t,i}} := \mathbb{D}_{t,x} \times \mathbb{D}_{t,y}$ around each $P_{t,i}$ such that $\mathcal{C} \cap \partial \mathbb{B}_{P_{t,i}} \subset \partial \mathbb{D}_{t,x} \times \mathbb{D}_{t,y}$ (we need condition (1.4) to ensure this). Let us denote by $\mu_{t,i}$ the local monodromy around each $P_{t,i}$ based at t_ε (close enough to t). Note that $\mu_{t,i} \in \mathbb{B}_m$, where $m = 1$ if and only if $P_{t,i}$ is a transversal intersection of \mathcal{C} and L_t . If $P_{t,i}$ is singular in \mathcal{C} , then the link produced by closing $\mu_{t,i}$ is the link of the singularity at $P_{t,i}$. If $P_{t,i}$ is a single tangency, then $\mu_{t,i}$ is a half twist. Therefore, $\mu_{t,i}$ is trivial if and only if $P_{t,i}$ is a transversal intersection of \mathcal{C} and L_t . We disregard these trivial braids, denote by κ_t the number of non-trivial braids, and denote them by $\mu_{t,1}, \dots, \mu_{t,\kappa_t}$ after reordering. Define by β_t a path that joins t_ε and t consider $\tilde{\beta}_t$ the open braid associated with β_t and define $\tau_{t,i} := \beta_t \cdot \mu_{t,i} \cdot \beta_t^{-1}$ (here “ \cdot ” just means juxtaposition). Since the strings inside each Milnor polydisk are different, the braids $\tau_{t,i}$ automatically commute.

For the *moreover* part, note that moving P_y generically on L_∞ causes just a small change of coordinates $X \mapsto X + \eta Y$ and hence each L_t splits into κ_t non-transversal vertical lines satisfying the genericity condition (1.4) at only one point. This shows the statement. \square

4.3. From generic at infinity to locally generic.

In the more general case when triples $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ are just generic at infinity, the change of projection might lead to more complicated situations, but in general the result is of the same nature, that is, some braids produced by the monodromy around \bar{L}_t in the non-generic braid monodromy need to be replaced by an appropriate factorization. We state some interesting particular cases.

Proposition 4.4. *Let us assume that L_t intersects $\bar{\mathcal{C}}$ transversally at $k - 3$ points and it is tangent to an ordinary cusp. After conjugation the monodromy around L_t produces the braid $\mu := (\sigma_2\sigma_1)^2$.*

Moreover, μ can be replaced by the factorization $(\sigma_1^{\sigma_2}, \sigma_2^3)$ so that the monodromy becomes generic in a regular neighborhood of L_t .




FIGURE 7. Changing the projection.

Proof. The local equation of \mathcal{C} at P is $y^3 - x^2 = 0$. Note that the local monodromy around $t = 0$ is given by the braid parametrized by $y^3 = e^{4\pi\sqrt{-1}\lambda}$, $\lambda \in [0, 1]$, that is, a rotation of angle $\frac{4}{3}\pi$ of the 3-rd roots of unity. This produces the braid $(\sigma_2\sigma_1)^2$. A small perturbation of the projection point along \bar{L}_∞ produces a change of variable $f_\eta := y^3 - (x - 3\eta y)^2$. The discriminant of f_η with respect to y is $x^3(x - 4\eta^3)$, which implies that the non-transversal vertical line \bar{L}_0 splits into \bar{L}_0 and \bar{L}_{t_1} , where $t_1 = 4\eta^3$. It is straightforward to check that there is a cusp at $(0, 0)$ (whose tangent $x = 3\eta y$ is not vertical if $\eta \neq 0$), a simple point at $(0, 9\eta^2)$, a vertical tangency at $(t_1, 4\eta^2)$, and a simple point at (t_1, η^2) (see Figure 7).

Regardless of the value of η , there is a geometric basis on a local disk $\mathbb{D}_{x,0}$ centered at $t = 0$ and a choice of generators σ_1 and σ_2 in \mathbb{B}_3 such that the local braid monodromy around $t = 0$ is given by σ_2^3 . Therefore, since the product should be $(\sigma_2\sigma_1)^2$ and $\sigma_2^3 \cdot \sigma_1^{\sigma_2} = \sigma_2^2\sigma_1\sigma_2 = (\sigma_2\sigma_1)^2$, then the local monodromy around $t = t_1$ should be given by $\sigma_1^{\sigma_2}$, which concludes the proof. \square

Similar computations provide the following results for the simplest cases:

Proposition 4.5. *Let us assume that L_t intersects $\bar{\mathcal{C}}$ transversally at $k - 3$ points and it is tangent at one of the local branches of an ordinary double point. After conjugation the monodromy around L_t produces the braid $\mu := \sigma_1\sigma_2\sigma_1$.*

Moreover, μ can be replaced by the factorization $(\sigma_2^{\sigma_1}, \sigma_1^2)$ so that the monodromy becomes generic in a regular neighborhood of L_t .

Proposition 4.6. *Let us assume that L_t intersects $\bar{\mathcal{C}}$ transversally at $k - m$ points and it is tangent to an inflection point of order m . After conjugation the monodromy around L_t produces the braid $\mu := \sigma_{m-1} \dots \sigma_1$.*

Moreover, μ can be replaced by the factorization $(\sigma_1, \dots, \sigma_{m-1})$ so that the monodromy becomes generic in a regular neighborhood of L_t .

4.4. Line arrangements: from non-generic to generic.

We are also interested in the case where $\bar{\mathcal{C}}$ is the union of a fully horizontal curve $\bar{\mathcal{C}}_0$ (such that $P_y \notin \bar{\mathcal{C}}$) and some lines $\bar{L}_1, \dots, \bar{L}_\ell$ passing through P_y and different from \bar{L}_∞ . In order to avoid technical problems, we restrict our attention to the case of line arrangements, which admits two approaches. One of them is to use wiring diagrams [8] (which is an invariant equivalent to braid monodromy). The other is more direct but we need some definitions.

Definition 4.7. Let $n, k \in \mathbb{N}$ and let $\ell \in \{0, 1, \dots, k\}$. The ℓ -shift of \mathbb{B}_n into \mathbb{B}_{n+k} is the inclusion $\rho_\ell : \mathbb{B}_n \rightarrow \mathbb{B}_{n+k}$ such that $\rho(\sigma_j) = \sigma_{j+\ell}$, $1 \leq j < n$.

Definition 4.8. The *partial Garside element* of the strings i, \dots, j , $i \leq j$, is the image $\Delta_{i,j}$ of the Garside element by the $(i-1)$ -shift of \mathbb{B}_{j-i+1} into \mathbb{B}_n .

Let $\bar{\mathcal{C}} = \bar{L}_1 \cup \dots \cup \bar{L}_n \cup \bar{L}_{n+1} \cup \dots \cup \bar{L}_{n+k}$ be a line arrangement such that P_y is a point of multiplicity k and $P_y \in \bar{L}_{n+1} \cap \dots \cap \bar{L}_{n+k}$. Let \bar{L}_∞ be a generic line through P_y . We want to construct a generic braid monodromy for $\bar{\mathcal{C}}$ starting from the one of $(\bar{\mathcal{C}}_h, P_y, \bar{L}_\infty)$, where $\bar{\mathcal{C}}_h := \bar{L}_1 \cup \dots \cup \bar{L}_n$. Recall that this braid monodromy is obtained as a representation $\nabla : \pi_1(\mathbb{C} \setminus \mathcal{B}; t_0) \rightarrow \mathbb{B}_n$ where \mathcal{B} is the set of x -coordinates of the multiple points of \mathcal{C}_h . Let $\mathcal{B}_0 := \mathcal{B} \cup \{t_1, \dots, t_k\}$, where $L_{n+i} = \{x = t_i\}$; since the braids associated with the meridians around the points in $\mathcal{B}_0 \setminus \mathcal{B}$ are trivial, the above mapping defines a representation $\nabla_0 : \pi_1(\mathbb{C} \setminus \mathcal{B}_0; t_0) \rightarrow \mathbb{B}_n$ (which will be referred to as the *augmented braid monodromy*). A choice of an ordered geometric basis $(\gamma_1, \dots, \gamma_r)$ allows to represent this braid monodromy by $(\tau_1, \dots, \tau_r) \in \mathbb{B}_n^r$; let $1 \leq i_1 < \dots < i_k \leq r$ be the indices of the braids corresponding to the vertical lines L_{n+1}, \dots, L_{n+k} .

Proposition 4.9. *Under the above conditions, let us decompose τ_i as $\beta_i * \alpha_i$ where*

$$(4.1) \quad \alpha_i = \prod_{s=1}^{m(i)} \Delta_{a_s(i), a_{s+1}(i)-1}^2, \quad 1 = a_1(i) < \dots < a_{m(i)+1}(i) = n+1,$$

is a product of squares of partial Garside elements of \mathbb{B}_n . Then, a generic braid monodromy for $\bar{\mathcal{C}}$ is obtained by replacing the braids τ_i as follows:

- (1) *If $i_j < i < i_{j+1}$ (by convention $i_0 = 0$, $i_{k+1} = r+1$) then replace τ_i by the sequence*

$$\left\{ \beta_i * \Delta_{a_s(i), a_{s+1}(i)-1}^2 \right\}_{m(i) \geq s \geq 1}$$

and we take out the trivial braids (for s such that $a_s(i) < a_{s+1}(i) - 1$).

- (2) *If $i = i_j$ then replace τ_i by*

$$\{(\beta_i \cdot \sigma_{n+j-1}^{-1} \cdot \dots \cdot \sigma_{n+1}^{-1} \cdot \Delta_{a_{m(i)}(i), a_{m(i)+1}(i)} \cdot \dots \cdot \Delta_{a_{s+1}(i), a_{s+2}(i)}) * \Delta_{a_s(i), a_{s+1}(i)}^2\}_{m(i) \geq s \geq 1}.$$

- (3) *Finally, add $\Delta_{n+1, n+k}^2$.*

Proof. For simplicity, we can assume that $r = k$, since we may add vertical lines to the arrangement and once the generic braid monodromy is obtained we may forget the strings corresponding to the added lines; it is easily seen that this does not affect to the final result.

The augmented braid monodromy ∇_0 described above may be computed as follows. Fix a generic line \bar{L}_* through P_y close to \bar{L}_∞ and fix a base point $P_* := (t_0, y_*) \in L_*$ close to P_y . We fix an ordered geometric basis (μ_1, \dots, μ_n) of $\pi_1(L_* \setminus \mathcal{C}_h; P_*)$ such that μ_i is a meridian of L_i and its pseudo-Coxeter element is the the boundary of a disk \mathbb{D}_* in L_* surrounding $L_* \cap \mathcal{C}_h$ (which is also the negative boundary of a disk in \bar{L}_* centered at P_y).

In the affine plane $\mathbb{C}^2 = \mathbb{P}^2 \setminus \bar{L}_\infty$, we consider the horizontal line H_* passing through P_* . We can choose P_* such that there is a disk \mathbb{D}_x containing $\mathcal{B}_0 \times \{y_*\}$ in its interior. We fix an ordered geometric basis $(\tilde{\mu}_1, \dots, \tilde{\mu}_n, \tilde{\gamma}_1, \dots, \tilde{\gamma}_k)$ of $\pi_1(H_* \setminus \mathcal{C}; P_*)$ such that:

- The meridians $\tilde{\mu}_i$ and μ_i are equal in $\pi_1(\mathbb{C}^2 \setminus \mathcal{C}; P_*)$ (and we identify them from now on). The product $\tilde{\mu}_n \dots \tilde{\mu}_1$ is the boundary of a disk $\tilde{\mathbb{D}}_*$ which is isotopic to \mathbb{D}_* (in $\mathbb{C}^2 \setminus \mathcal{C}$) and disjoint to \mathbb{D}_x (see Figure 8(a)).
- The meridians $(\tilde{\gamma}_1, \dots, \tilde{\gamma}_k)$ are obtained as follows. Pick a base point in $\partial\mathbb{D}_x$; all the meridians have as common part a path joining P_* with this point avoiding (counterclockwise) the disk $\tilde{\mathbb{D}}_*$. These meridians project onto the meridians γ_i .

It is then clear that $(\tilde{\gamma}_k \dots \tilde{\gamma}_1) \cdot (\mu_d \dots \mu_1) = 1$ in $\pi_1(\mathbb{P}^2 \setminus \bar{\mathcal{C}}; P_*)$.

Next step is to deform the projection point in \bar{L}_∞ which induces a family of coordinate changes $(x, y) \mapsto (x + s\varepsilon y, y)$ for a fixed $0 < |\varepsilon| \ll 1$ and $s \in (0, 1]$. For each new projection point we obtain a braid monodromy $\nabla_s : \pi_1(\mathbb{C} \setminus \mathcal{B}_s; t_{0,s}) \rightarrow \mathbb{B}_{n+k}$, $s \in (0, 1]$. We can fix small pairwise disjoint disks $\mathbb{D}_1, \dots, \mathbb{D}_k$ (respectively centered at t_1, \dots, t_k) such that \mathcal{B}_s is contained in the interior of $\bigcup_{j=1}^k \mathbb{D}_j$. Note that $\#(\mathcal{B}_s \cap \mathbb{D}_j) = \#(\mathcal{C}_h \cap L_{n+j}) = m(j)$. In order to compute the braid monodromy ∇_s we choose an ordered geometric basis $(\eta_{i,j})_{1 \leq j \leq k, 1 \leq i \leq m(j)}$, such that $\prod_{i=1}^{m(j)} \eta_{i,j} = \gamma_j$. The generic base fiber is the new vertical line passing through P_* . In this vertical line we can construct disks isotopic to $\tilde{\mathbb{D}}_*$ and \mathbb{D}_x containing respectively the intersections with $L_1 \cup \dots \cup L_n$ and $L_{n+1} \cup \dots \cup L_{n+k}$.

We start with the computation of $\nabla_1(\gamma_j)$. The decomposition $\tau_j = \beta_j * \alpha_j$ has a geometric meaning: β_j is the braid starting from the vertical line through P_* and ending in a vertical line close to L_{n+j} while α_j is the braid obtained by turning around L_{n+j} as (4.1) justifies. Let us produce a similar decomposition for $\nabla_1(\gamma_j) = \tilde{\beta}_j * \tilde{\alpha}_j$ in \mathbb{B}_{n+k} . The motion producing β_j induces now a braid $\tilde{\beta}_j$ as the product of two commuting braids: β_j and $\sigma_{n+j-1}^{-1} \dots \sigma_{n+1}^{-1}$, see Figure 8(b). The first one corresponds to the motion of the points in the disk $\tilde{\mathbb{D}}_*$ and the second one is obtained by considering the motion of the i^{th} point to the boundary (behind the other ones), and then how this point approaches to $\tilde{\mathbb{D}}_*$.

The braid $\tilde{\alpha}_j$ is decomposed as a product of conjugate of partial Garside elements leading to the images of $\eta_{i,j}$ as in (2), see Figure 9.

For the case $k < r$, the decomposition in (1) is obtained is the same one by forgetting the strings corresponding to the verticals line not in \mathcal{C} .





FIGURE 8.

FIGURE 9. Braid $\tilde{\alpha}_j$.

The final step consists in the choice of a new line at infinity close to \bar{L}_∞ (without changing the projection point); the last lines are no more parallel since the multiple point P_y becomes affine and we must add the braid of (3). \square

Example 4.10. Let us consider the line arrangement of Figure 10. In order to obtain the augmented braid monodromy ∇_0 it is necessary to add an extra vertical line L passing through the double point $L_1 \cap L_3$ of \mathcal{C} . It is easy to see that ∇_0 is given by $(\sigma_2^2, 1, \sigma_2 * \sigma_1^2, \sigma_1^2)$. The generic braid monodromy is given by:

$$(\Delta_{2,4}^2, \Delta_{2,4} * \sigma_1^2, (\sigma_3^2)^{\sigma_4}, (\sigma_4^{-1} \sigma_3) * \sigma_2^2, (\sigma_4^{-1} \sigma_3 \sigma_2) * \sigma_1^2, \sigma_2 * \sigma_1^2, (\sigma_3^2)^{\sigma_4 \sigma_5}, (\sigma_5^{-1} \sigma_4^{-1} \sigma_3) * \Delta_{1,3}^2).$$




FIGURE 10. From non-generic to generic braid monodromy.

5. TRANSFORMATIONS OF CURVES BY KUMMER COVERS.

Let \bar{C} be a (reduced) projective curve of degree k of equation $F_k(x, y, z) = 0$ and let \bar{C}_n be its transform by a Kummer cover π_n , $n > 1$. Note that \bar{C}_n is a projective curve of degree nk of equation $F_k(x^n, y^n, z^n) = 0$. Another obvious remark is that if \bar{C} is reducible so is \bar{C}_n . The converse is not true as we will see in §6.

We will briefly analyze the singularities of \bar{C}_n in terms of \bar{C} . For convenience we distinguish three types of points in \mathbb{P}^2 .

Definition 5.1. Let $P \in \mathbb{P}^2$ such that $P := [x_0 : y_0 : z_0]$. We say that P is a point of *type* $(\mathbb{C}^*)^2$ (or simply of *type* 2) if $x_0 y_0 z_0 \neq 0$. If $x_0 = 0$ but $y_0 z_0 \neq 0$ the point is said to be of *type* \mathbb{C}_x^* (types \mathbb{C}_y^* and \mathbb{C}_z^* are defined accordingly). Such points will also be referred to as *type* 1 points. The corresponding line (either $L_X := \{X = 0\}$, $L_Y := \{Y = 0\}$, or $L_Z := \{Z = 0\}$) a type-1 lies on will be referred to as their *axis*. The remaining points $P_x := [1 : 0 : 0]$, $P_y := [0 : 1 : 0]$, and $P_z := [0 : 0 : 1]$ will be called *vertices* (or type 0 points) and their axes are the two lines (either L_X , L_Y , or L_Z) they lie on.

Remark 5.2. Note that a point of type ℓ , $\ell = 0, 1, 2$ in \mathbb{P}^2 has exactly n^ℓ preimages under π_n . It is also clear that the local type of \bar{C}_n at any two points on the same fiber are analytically equivalent.

Lemma 5.3. Let $P \in \mathbb{P}^2$ be a point of type ℓ and $Q \in \pi_n^{-1}(P)$. There exist local coordinates (u_0, v_0) and (u_1, v_1) centered at Q and P , respectively, such that:

- (1) If $\ell = 2$, then $(u_1, v_1) = \pi_n(u_0, v_0) = (u_0, v_0)$.
- (2) If $\ell = 1$, then $(u_1, v_1) = \pi_n(u_0, v_0) = (u_0^n, v_0)$, where $u_0 = 0$ and $u_1 = 0$ are the local equations (at Q and P , respectively) of the axes containing the points.
- (3) If $\ell = 0$, then $(u_1, v_1) = \pi_n(u_0^n, v_0) = (u_0^n, v_0^n)$, where $u_i = 0$ and $v_i = 0$ are the local equations of the axes containing P, Q .

Proof. For the points of type 2, it is a straightforward consequence of fact that the cover π_n is unramified at P . If P is of type 1, let us assume that $Q := [0 : t : 1]$, $t \in \mathbb{C}^*$, and then

$P = [0 : t^n : 1]$. We choose $(u_0, v_0) = (x, y - t)$ and $(u_1, v_1) = (x, \sqrt[n]{t^n + y})$. If P is of type 0, we may assume $Q = [0 : 0 : 1]$; we keep x, y as coordinates for P and Q . \square

Proposition 5.4. *Let $P \in \mathbb{P}^2$ be a point of type ℓ and $Q \in \pi_n^{-1}(P)$. One has the following:*

- (1) *If $\ell = 2$, then $(\bar{\mathcal{C}}, P)$ and $(\bar{\mathcal{C}}_n, Q)$ are analytically isomorphic.*
- (2) *If $\ell = 1$, then $(\bar{\mathcal{C}}_n, Q)$ is a singular point of type 1 if and only if $m > 1$, where $m := (\bar{\mathcal{C}} \cdot \bar{L})_P$ and \bar{L} is the axis of P .*
- (3) *If $\ell = 0$, then $(\bar{\mathcal{C}}_n, Q)$ is a singular point.*

Proof. It is a straightforward consequence of Lemma 5.3. Assume $P = Q = (0, 0)$ are points of type ℓ on $\bar{\mathcal{C}}$ and $f(u_1, v_1)$ is a local equation of $(\bar{\mathcal{C}}, P)$. If $\ell = 2$, then $f(u_0, v_0)$ is a local equation of $(\bar{\mathcal{C}}_n, Q)$. If $\ell = 1$ then $f(u_0^n, v_0)$ is a local equation of $(\bar{\mathcal{C}}_n, Q)$. Finally, if $\ell = 0$, then $f(u_0^n, v_0^n)$ is a local equation of $(\bar{\mathcal{C}}_n, Q)$. \square

Example 5.5. In some cases, we can be more explicit about the singularity type of $(\bar{\mathcal{C}}_n, Q)$.

- (1) If P is of type 1, $(\bar{\mathcal{C}}, P)$ is smooth and $m := (\bar{\mathcal{C}} \cdot \bar{L})_P$ then $(\bar{\mathcal{C}}_n, Q)$ has the same topological type as $u_0^n - v_0^m = 0$. In particular, if $n = 2$, then $(\bar{\mathcal{C}}_n, Q)$ is of type \mathbb{A}_{m-1} .
- (2) If P is of type 0, $(\bar{\mathcal{C}}, P)$ is smooth and transverse to the axes then $(\bar{\mathcal{C}}_n, Q)$ is an ordinary multiple point of multiplicity n .

In order to better describe singular points of type 0 and 1 of $\bar{\mathcal{C}}_n$ we will introduce some notation. Let $P \in \mathbb{P}^2$ be a point of type $\ell = 0, 1$ and $Q \in \pi_n^{-1}(P)$ a singular point of $\bar{\mathcal{C}}_n$. Denote by μ_P (resp. μ_Q) the Milnor number of $\bar{\mathcal{C}}$ at P (resp. $\bar{\mathcal{C}}_n$ at Q), also, denote by $\delta_1, \dots, \delta_r$ the local branches of $\bar{\mathcal{C}}$ at P and consider $\tilde{\delta}_i := \pi_n^{-1}(\delta_i)$. Define μ_{P, δ_i} (resp. $\mu_{Q, \tilde{\delta}_i}$) as the Milnor number of the singularity (δ_i, P) (resp. $(\tilde{\delta}_i, Q)$). Since $\ell = 0, 1$, then P and Q belong to either exactly one or two axes. If P and Q belong to an axis \bar{L} , then $m_P^{\bar{L}} := (\bar{\mathcal{C}} \cdot \bar{L})_P$ and $m_{P, \delta_i}^{\bar{L}} := (\delta_i \cdot \bar{L})_P$ (analogous notation for Q and $\tilde{\delta}_i$). More specific details about singular points of types 0 and 1 can be described as follows:

Proposition 5.6. *Under the above conditions and notation one has the following:*

- (1) *For $\ell = 1$, P belongs to a unique axis \bar{L} and*
 - (a) $\mu_Q = n\mu_P + (m_P^{\bar{L}} - 1)(n - 1)$,
 - (b) $\mu_{Q, \tilde{\delta}_i} = n\mu_{P, \delta_i} + (m_{P, \delta_i}^{\bar{L}} - 1)(n - 1)$,
 - (c) *If $i \neq j$ then $(\tilde{\delta}_i \cdot \tilde{\delta}_j)_Q = n(\delta_i \cdot \delta_j)_P$,*
 - (d) *If $r_i := \gcd(n, m_{P, \delta_i}^{\bar{L}})$, then $\tilde{\delta}_i$ has r_i irreducible components.*
- (2) *For $\ell = 0$, P belongs to exactly two axes \bar{L}_1 and \bar{L}_2*
 - (a) $\mu_Q = n^2\mu_P + (n - 1)(n(m_P^{\bar{L}_1} + m_P^{\bar{L}_2}) - 1)$,
 - (b) $\mu_{Q, \tilde{\delta}_i} = n^2\mu_{P, \delta_i} + (n - 1)(n(m_{P, \delta_i}^{\bar{L}_1} + m_{P, \delta_i}^{\bar{L}_2}) - 1)$, $m_{Q, \tilde{\delta}_i}^{\bar{L}_j} = nm_{P, \delta_i}^{\bar{L}_j}$,
 - (c) *If $i \neq j$ then $(\tilde{\delta}_i \cdot \tilde{\delta}_j)_Q = n^2(\delta_i \cdot \delta_j)_P$,*
 - (d) *If $r_i := \gcd(n, m_{P, \delta_i}^{\bar{L}_1}, m_{P, \delta_i}^{\bar{L}_2})$, then $\tilde{\delta}_i$ has nr_i irreducible components which are analytically isomorphic to each other for any fixed i .*

Proof. For part ((1)) note that

$$\begin{aligned}\mu_Q &= \dim_{\mathbb{C}} \frac{\mathbb{C}\{x,y\}}{(x^{n-1}f_x(x^n,y), f_y(x^n,y))} = \dim_{\mathbb{C}} \frac{\mathbb{C}\{x,y\}}{(x^{n-1}, f_y(x,y))} + \dim_{\mathbb{C}} \frac{\mathbb{C}\{x,y\}}{(f_x(x^n,y), f_y(x^n,y))} = \\ &= (n-1) \dim_{\mathbb{C}} \frac{\mathbb{C}\{x,y\}}{(x, f_y(x,y))} + n \dim_{\mathbb{C}} \frac{\mathbb{C}\{x^n,y\}}{(f_x(x^n,y), f_y(x^n,y))} = (n-1)(m_P^{\bar{L}} - 1) + n\mu_P.\end{aligned}$$

The same proof applies to $\mu_{Q, \tilde{\delta}_i}$. Also

$$(\tilde{\delta}_i \cdot \tilde{\delta}_j)_Q = \dim_{\mathbb{C}} \frac{\mathbb{C}\{x,y\}}{(f_i(x^n,y), f_j(x^n,y))} = n \dim_{\mathbb{C}} \frac{\mathbb{C}\{x^n,y\}}{(f_i(x^n,y), f_j(x^n,y))} = n(\delta_i \cdot \delta_j)_P,$$

where f_i (resp. f_j) is a local equation for δ_i (resp. δ_j).

Finally, we can describe the irreducible branch δ_i as a Puiseux factorization of type $f_i(x,y) = \prod_{j=1}^{\nu} (y - s(\xi_{\nu}^j x^{1/\nu}))$, where ν is a multiple of $m = m_{P, \delta_i}^{\bar{L}}$, the order of f_i in y . Note that $f_i(x^n, y) = \prod_{j=1}^{\nu} (y - s(\xi_{\nu}^j x^{n/\nu}))$. Assume r is a common divisor, that is, $n = rn'$, $m = rm'$ (and hence $\nu = r\nu'$) then

$$\prod_{j=1}^{\nu'} (y - s(\xi_{\nu'}^{rj} x^{rn'/\nu'})) = \prod_{j=1}^{\nu'} (y - s(\xi_{\nu'}^j x^{n'/\nu'}))$$

is invariant under the Galois conjugation by ν' -roots of unity and hence it is a convergent series in $\mathbb{C}\{x,y\}$, that is, a union of branches. Therefore the result follows.

The proof of part (2) is analogous. \square

6. EXAMPLES AND APPLICATIONS.

In this section the results obtained in §2-3 will be applied to produce generic braid monodromies of curves. We will follow the conventions introduced in [5]. In the language of the previous section, diagram systems on \mathbb{C} for the starting curve \mathcal{C} will be chosen by joining the points with segments in a decreasing order according to the lexicographic order.

6.1. Smooth curves.

We start with the braid monodromy of a smooth curve considered as a Fermat curve $\bar{\mathcal{C}}_n = \{x^n - y^n + z^n = 0\} \subset \mathbb{P}^2$, via the n -th Kummer cover of the line $\bar{\mathcal{C}}$ of equation $x - y + z = 0$. Using the conventions of §2, one can obtain an extended braid monodromy $\tilde{\nabla} : \pi_1(\mathbb{C} \setminus \tilde{B}; t_0) \rightarrow \mathbb{B}_{1,1}$, where γ_1, γ_2 is a geometric basis of the free group $\pi_1(\mathbb{C} \setminus \tilde{B}; t_0)$ and $\tilde{\nabla}(\gamma_1) = 1$, $\tilde{\nabla}(\gamma_2) = \sigma_1^2$ as shown in Figure 11. Using Example 3.5 and the commutative diagram (2.2), one can obtain a braid monodromy $\nabla_n : \mathbb{F}_n \rightarrow \mathbb{B}_n$ for \mathcal{C}_n as follows:

$$(6.1) \quad \nabla_n(\tilde{\gamma}_j) = \sigma_{n-1} \cdot \dots \cdot \sigma_1,$$

where the basis $\tilde{\gamma}_j$, $j = 1, \dots, n$ is obtained as in Lemma 2.7, by forgetting the last term since \bar{L}_0 is transversal.

However, note that ∇_n is not a generic braid monodromy since the projection from P_y contains n special fibers $\bar{L}_{\xi_n^i}$, $i = 0, \dots, n-1$ with an order n tangency at $[\xi_n^i : 0 : -\xi_n^i]$. Still, ∇_n is useful to compute the fundamental group of the complement of a smooth curve (using Theorem 1.6). This group is generated by μ_1, \dots, μ_d with a relation $\mu_d \cdot \dots \cdot \mu_1$ and since

$$\mu_i^{\sigma_{n-1} \cdot \dots \cdot \sigma_1} = \mu_{i+1}, \quad 1 \leq i < d$$




FIGURE 11. A generic line.

the group is cyclic of order d , as obtained by Zariski in [35].

By the previous discussion, in order to obtain a generic braid monodromy, it is enough to use Proposition 4.6 and obtain the following.

Proposition 6.1 ([30, Theorem 1, p. 120]). *Let \bar{C} be a smooth curve of degree n . Then \bar{C} induces a braid monodromy factorization of Δ_n^2 given by*

$$\overbrace{(\sigma_1, \dots, \sigma_{n-1})}^{1^{st} \text{ package}}, \dots, \overbrace{(\sigma_1, \dots, \sigma_{n-1})}^{n^{th} \text{ package}}.$$

6.2. Zariski sextics.

In [35] Zariski showed that the fundamental group of a sextic with six cusps on a conic is $\mathbb{Z}/2 * \mathbb{Z}/3$ and the family of such sextics is irreducible, so they all have equivalent braid monodromies. We are going to compute one in two steps: first a (fully horizontal) non-generic braid monodromy of such a sextic will be obtained from a simpler curve via a Kummer covering and second, a deformation will be performed to compute the desired generic braid monodromy.

Consider a conic as in Figure 12. It is tangent to two of the axes and transversal to the third one, e.g., the conic $\bar{C} = \{x^2 + y^2 - 2xz - 2yz + z^2 = 0\}$. Using the conventions of §2, the projection from P_y has three non-generic fibers at $\tilde{B} = \{0, 1, 2\}$. Consider $t_0 = 3$ and $\gamma_1, \gamma_2, \gamma_3$ meridians around 2, 1, and 0 resp. (see Figure 12) forming an ordered geometric basis of the free group $\pi_1(\mathbb{C} \setminus \tilde{B}; t_0)$. One can obtain an extended braid monodromy $\tilde{\nabla} : \pi_1(\mathbb{C} \setminus \tilde{B}; t_0) \rightarrow \mathbb{B}_{2,1}$ as follows

$$\tilde{\nabla}(\gamma_1) = \sigma_1, \quad \tilde{\nabla}(\gamma_2) = \sigma_1 * \sigma_2^4, \quad \tilde{\nabla}(\gamma_3) = (\sigma_1 \sigma_2^2) * \sigma_1.$$

After conjugating by the braid $(\sigma_1 \sigma_2^2)$, the above representation becomes:

$$\tilde{\nabla}(\gamma_1) = \sigma_1^2, \quad \tilde{\nabla}(\gamma_2) = \sigma_2^4, \quad \tilde{\nabla}(\gamma_3) = \sigma_1.$$

Let us consider the 3-rd Kummer cover of \bar{C} . The curve \bar{C}_3 is a curve of degree 6. Since \bar{C} is smooth and intersects transversally \bar{L}_∞ , the singular points are contained in $\bar{L}_0 \cup \bar{L}_Y$ (Proposition 5.4). Using Proposition 5.6, the curve \bar{C}_3 possesses six ordinary cusps (which are in a conic, namely the union of the lines $\pi_3^{-1}(\bar{L}_0) = \bar{L}_0$ and $\pi_3^{-1}(\bar{L}_Y) = \bar{L}_Y$).




FIGURE 12. A conic tangent to two axes.

Using a radial system of generators as in §3.2, the map $\hat{\rho}_3 : \mathbb{B}_{2,1} \rightarrow \mathbb{B}_6$ is given (see Proposition 3.7) by:

$$\begin{aligned}\sigma_1 &\mapsto \sigma_1 \sigma_3 \sigma_5 \\ \sigma_2^2 &\mapsto \sigma_5 \sigma_4 \sigma_3 \sigma_2 \sigma_3^{-1} \sigma_5^{-1} = (\sigma_5 \sigma_4 \sigma_2^{-1}) * (\sigma_4 \sigma_3),\end{aligned}$$

where the generators σ_i on the right-hand side are defined as $\sigma_1 := \tilde{\sigma}_{1,1}, \sigma_2 := \tilde{\sigma}_{2,1}, \dots, \sigma_5 := \tilde{\sigma}_{1,3}$ (with the right-lexicographic order) for simplicity. Let us conjugate the result by $\sigma_5 \sigma_3$:

$$(6.2) \quad \begin{aligned}\sigma_1 &\mapsto \sigma_1 \sigma_3 \sigma_5 \\ \sigma_2^2 &\mapsto \sigma_3^{-1} \sigma_4 \sigma_3 \sigma_2 = \sigma_4 * (\sigma_3 \sigma_2).\end{aligned}$$

Proposition 6.2. *There is a geometric basis, as in Lemma 2.7, such that $\nabla_3 : \mathbb{F}_7 \rightarrow \mathbb{B}_6$ is given as follows:*

$$\begin{aligned}\tilde{\gamma}_1 &= \gamma_1 \mapsto (\sigma_1 \sigma_3 \sigma_5)^{\sigma_4 * (\sigma_3 \sigma_2)} = (\sigma_1 \sigma_3 \sigma_5)^{\sigma_4 \sigma_3 \sigma_2} \\ \tilde{\gamma}_2 &= \gamma_2 \mapsto \sigma_4 * (\sigma_3 \sigma_2)^2 \\ \tilde{\gamma}_3 &= \gamma_1^{\gamma_3} \mapsto (\sigma_1 \sigma_3 \sigma_5)^{\sigma_4 * (\sigma_3 \sigma_2) \sigma_1 \sigma_3 \sigma_5} = (\sigma_1 \sigma_3 \sigma_5)^{\sigma_4 \sigma_3 \sigma_2 \sigma_1 \sigma_3 \sigma_5} \\ \tilde{\gamma}_4 &= \gamma_2^{\gamma_3} \mapsto ((\sigma_3 \sigma_2)^2)^{\sigma_4^{-1} \sigma_1 \sigma_3 \sigma_5} \\ \tilde{\gamma}_5 &= \gamma_1^{\gamma_3^2} \mapsto (\sigma_1 \sigma_3 \sigma_5)^{\sigma_4 * (\sigma_3 \sigma_2) \sigma_1^2 \sigma_3^2 \sigma_5^2} = (\sigma_1 \sigma_3 \sigma_5)^{\sigma_4 \sigma_3 \sigma_2 \sigma_1^2 \sigma_3^2 \sigma_5^2} \\ \tilde{\gamma}_6 &= \gamma_2^{\gamma_3^2} \mapsto ((\sigma_3 \sigma_2)^2)^{\sigma_4^{-1} \sigma_1^2 \sigma_3^2 \sigma_5^2} \\ \tilde{\gamma}_7 &= \gamma_3^3 \mapsto \sigma_1^3 \sigma_3^3 \sigma_5^3\end{aligned}$$

As mentioned above, this braid monodromy is not generic. However, the triple $(\bar{\mathcal{C}}_3, P_y, L_\infty)$ is fully horizontal and transversal at infinity. One can deform ∇_3 as described in §4. First, a deformation is performed to obtain as in §4.3 to produce a locally generic triple. Then, a move as described in §4.2 will produce a generic braid monodromy.

Also note that the non-generic fibers of the projection from P_y are

$$\tilde{B}_3 = \{t_1 = \sqrt[3]{2}, t_2 = 1, t_3 = \xi_3 \sqrt[3]{2}, t_4 = \xi_3, t_5 = \xi_3^2 \sqrt[3]{2}, t_6 = \xi_3^2, t_7 = 0\},$$

where $\tilde{\gamma}_i$ is a meridian around t_i and:

- (1) \bar{L}_{t_i} , $i = 1, 3, 5$ correspond to vertical lines containing three simple tangencies (hence locally generic fibers).
- (2) \bar{L}_{t_i} , $i = 2, 4, 6$ correspond to vertical lines passing tangent to an ordinary cusp and transversal to three smooth points of $\bar{\mathcal{C}}_3$.

- (3) \bar{L}_0 corresponds to a vertical line passing through three ordinary cusps (hence a locally generic fiber).

Step 1. This step is necessary at the fibers of type (2) above. According to Proposition 4.4 the braid obtained as image of $\tilde{\gamma}_2$, that is, $\sigma_4 * (\sigma_3 \sigma_2)^2 = ((\sigma_4 * \sigma_3) \cdot (\sigma_2))^2$ has to be replaced by

$$(6.3) \quad (\sigma_2^{\sigma_4 * \sigma_3}, \sigma_4 * \sigma_3^3) = (\sigma_2^{\sigma_3 \sigma_4^{-1}}, (\sigma_3^3)^{\sigma_4^{-1}}).$$

Analogously can be done with $\nabla_3(\tilde{\gamma}_4)$ and $\nabla_3(\tilde{\gamma}_6)$.

Step 2. This is necessary at the locally generic fibers described above. Applying Proposition 4.3 to the fibers of types (1) and (3) one obtains that, for instance, $\nabla_3(\tilde{\gamma}_1) = (\sigma_1 \sigma_3 \sigma_5)^{\sigma_4 \sigma_3 \sigma_2}$ is replaced by

$$(6.4) \quad (\sigma_1^{\sigma_4 \sigma_3 \sigma_2}, \sigma_3^{\sigma_4 \sigma_3 \sigma_2}, \sigma_5^{\sigma_4 \sigma_3 \sigma_2}) = (\sigma_1^{\sigma_2}, \sigma_4, \sigma_5^{\sigma_4 \sigma_3 \sigma_2}).$$

Analogously can be done with $\nabla_3(\tilde{\gamma}_3)$, $\nabla_3(\tilde{\gamma}_5)$, and $\nabla_3(\tilde{\gamma}_7)$.

However, instead of working out each $\nabla_3(\tilde{\gamma}_i)$ separately we can note that $\nabla_3(\tilde{\gamma}_{i+2j}) = \nabla_3(\tilde{\gamma}_i)^{\sigma_1^j \sigma_3^j \sigma_5^j}$, ($i = 1, 2, j = 0, 1, 2$). Therefore one can also work out the first set of braids coming from $\nabla_3(\tilde{\gamma}_1)$ and $\nabla_3(\tilde{\gamma}_2)$ and conjugate to obtain the rest of the braids coming from $\nabla_3(\tilde{\gamma}_i)$ $i = 3, \dots, 6$. Finally attach the braids $(\sigma_1^3, \sigma_3^3, \sigma_5^3)$ obtained from $\nabla_3(\tilde{\gamma}_7)$. From (6.3) and (6.4) one obtains

$$(\sigma_1^{\sigma_2}, \sigma_4, \sigma_5^{\sigma_4 \sigma_3 \sigma_2}, \sigma_2^{\sigma_3 \sigma_4^{-1}}, (\sigma_3^3)^{\sigma_4^{-1}}).$$

Using the sequence of Hurwitz moves $h_2^{-1} h_3^{-1} h_4^{-1} h_2 h_3$ (see (1.3)) one obtains the first block:

$$(6.5) \quad (\sigma_1^{\sigma_2}, \sigma_2^{\sigma_3}, \sigma_3^3, \sigma_4^{\sigma_5^{-1} \sigma_3^{-1}}, \sigma_4).$$

After conjugating (6.5) by $\sigma_1 \sigma_3 \sigma_5$ and applying h_2 one obtains the second block:

$$(6.6) \quad (\sigma_2^{\sigma_3}, \sigma_3^3, \sigma_2^{\sigma_3^{-1} \sigma_1}, \sigma_4, \sigma_4^{\sigma_3 \sigma_5}).$$

The third block is obtained from (6.6) using conjugation by $\sigma_1 \sigma_3 \sigma_5$ and applying $h_1 h_4^{-1}$:

$$(6.7) \quad (\sigma_3^3, \sigma_2^{\sigma_3^{-1} \sigma_1}, \sigma_2^{\sigma_1^2}, \sigma_4, \sigma_4^{\sigma_3 \sigma_5}).$$

Proposition 6.3. *A generic braid monodromy for $\bar{\mathcal{C}}_3$ is given by:*

$$(\sigma_1^{\sigma_2}, \sigma_2^{\sigma_3}, \sigma_3^3, \sigma_4^{\sigma_5^{-1} \sigma_3^{-1}}, \sigma_4, \sigma_2^{\sigma_3}, \sigma_3^3, \sigma_2^{\sigma_3^{-1} \sigma_1}, \sigma_4, \sigma_4^{\sigma_3 \sigma_5}, \sigma_3^3, \sigma_2^{\sigma_3^{-1} \sigma_1}, \sigma_1^3, \sigma_1^{\sigma_2}, \sigma_4, \sigma_4^{\sigma_3 \sigma_5}, \sigma_3^3, \sigma_5^3).$$

Remark 6.4. This result is also obtained in [30, Theorem 1(3), p. 160]. It is straightforward to compute the fundamental group of the complement of $\bar{\mathcal{C}}_3$ and to retrieve Zariski's computation. Moreover, this braid monodromy allows us to compute the homotopy type of $\mathbb{C}^2 \setminus \mathcal{C}_3$ (for a generic choice of line at infinity) using the method in [27]. It is easy to see that

$$\mathbb{C}^2 \setminus \mathcal{C}_3 \simeq (\mathbb{S}^3 \setminus \{\text{trefoil knot}\}) \vee \bigvee_{i=1}^{13} \mathbb{S}^2.$$

6.3. Dual of a smooth cubic.

The dual of a smooth cubic is a sextic with 9 cusps. Kummer covers allow to recover one of these curves easily.

Consider a conic $\bar{\mathcal{C}} := \{x^2 + y^2 + z^2 - 2(xy + xz + yz) = 0\}$ as in Figure 13. Projecting from $P_y = [0 : 1 : 0]$ as usual, one obtains two non-generic fibers \bar{L}_∞ and \bar{L}_0 with tangencies $P_3 := [1 : 1 : 0]$ and $P_1 := [0 : 1 : 1]$ respectively. Also note that \mathcal{C} is tangent to \bar{L}_Y at $P_2 := [1 : 0 : 1]$. Using the conventions of §2, an extended braid monodromy $\tilde{\nabla} : \pi_1(\mathbb{C} \setminus \tilde{B}; t_0) \rightarrow \mathbb{B}_{2,1}$, where $\tilde{B} = \{P_1, P_2, P_3\}$, γ_1, γ_2 are meridians around P_2 and P_1 respectively forming an ordered basis of the free group $\pi_1(\mathbb{C} \setminus \tilde{B}; t_0)$ and $\tilde{\nabla}(\gamma_1) = \sigma_2^4$, $\tilde{\nabla}(\gamma_2) = \sigma_2^2 * \sigma_1$. For simplicity, we conjugate this monodromy by σ_2^2 and obtain

$$(6.8) \quad \gamma_1 \mapsto \sigma_2^4, \quad \gamma_2 \mapsto \sigma_1.$$




FIGURE 13. Tritangent conic.

The preimage of $\bar{\mathcal{C}}$ by the Kummer cover of order 3 is a sextic $\bar{\mathcal{C}}_3$ with nine cusps as it can be deduced from Example 5.5. Let us compute the braid monodromy $(\bar{\mathcal{C}}_3, P_y, \bar{L}_\infty)$. Following the ideas in §6.2 one immediately obtains

$$\begin{aligned} \gamma_1 &\mapsto ((\sigma_3 \sigma_2)^2)^{\sigma_4^{-1}} \\ \gamma_1^3 &\mapsto ((\sigma_3 \sigma_2)^2)^{\sigma_4^{-1} \sigma_1 \sigma_3 \sigma_5} \\ \gamma_1^2 &\mapsto ((\sigma_3 \sigma_2)^2)^{\sigma_4^{-1} \sigma_1^2 \sigma_3^2 \sigma_5^2} \\ \gamma_3^3 &\mapsto \sigma_1^3 \sigma_3^3 \sigma_5^3. \end{aligned}$$

Since the line \bar{L}_∞ is not generic, one can apply Proposition 4.2 to obtain a braid monodromy that is generic at infinity. The pseudo-Coxeter element for the monodromy (6.8) is $c := \sigma_1 \sigma_2^4$ and hence $\Delta_3^6 c^{-3} = (\Delta_3^2 c^{-1})^3 = (\sigma_1^3)^{\sigma_2^2} \in \mathbb{B}_{2,1}$, whose image in \mathbb{B}_6 via $\hat{\rho}_3$ (see (6.2)) is $(\sigma_1^3 \sigma_3^3 \sigma_5^3)^{\sigma_4 \sigma_3 \sigma_2 \sigma_4^{-1}}$. Therefore

$$(((\sigma_3 \sigma_2)^2)^{\sigma_4^{-1}}, ((\sigma_3 \sigma_2)^2)^{\sigma_4^{-1} \sigma_1 \sigma_3 \sigma_5}, ((\sigma_3 \sigma_2)^2)^{\sigma_4^{-1} \sigma_1^2 \sigma_3^2 \sigma_5^2}, \sigma_1^3 \sigma_3^3 \sigma_5^3, (\sigma_1^3 \sigma_3^3 \sigma_5^3)^{\sigma_4 \sigma_3 \sigma_2 \sigma_4^{-1}})$$

is a braid monodromy factorization of Δ_6^2 generic at infinity for the Zariski sextic $\bar{\mathcal{C}}_3$. Finally, in order to obtain a generic braid monodromy one needs to apply Propositions 4.2 and 4.3 to slightly turn the projection generic and perform Hurwitz moves to simplify the braids. One can check that the final generic braid monodromy of $\bar{\mathcal{C}}_3$ is:

$$\sigma_2^3, \sigma_2^{\sigma_3 \sigma_4^{-1}}, \sigma_2^{\sigma_3 \sigma_4^{-1} \sigma_1 \sigma_3 \sigma_5}, \sigma_3^3, (\sigma_3^3)^{\sigma_4 \sigma_5}, \sigma_5^3, \sigma_2^{\sigma_3^{-1} \sigma_4 \sigma_1^2 \sigma_5^{-1}}, (\sigma_5^3)^{\sigma_4}, \sigma_1^3, (\sigma_1^3)^{\sigma_2}, \sigma_4^3, (\sigma_5^3)^{\sigma_4 \sigma_3 \sigma_2}.$$

The fundamental group of this curve complement has been extensively studied by Zariski in [37]. He also used deformation arguments to recover the fundamental group studied in [35], see §6.2, as well as to study the fundamental group of sextics with six cusps not on a conic.

Similar arguments can be used to describe the generic braid monodromy of any curve $\bar{\mathcal{C}}_n$, n odd, which is an irreducible curve of degree $3n$ and $9n$ singularities of type \mathbb{A}_{n-1} (see [16]).

6.4. Ceva arrangement.

We are going to use the classical Ceva arrangement (six lines joining four points in general position) to find the braid monodromy of the 9-Ceva arrangement

$$\mathcal{C} := \{(x^3 - y^3)(y^3 - z^3)(x^3 - z^3) = 0\},$$

and it can be obtained as a Kummer covering of three concurrent lines in a classical Ceva arrangement, see Figure 14.




FIGURE 14. Ceva arrangement.

The extended braid monodromy on $\mathbb{B}_{2,1}$ of Figure 14 is given by:

$$\gamma_1 \mapsto \sigma_1^2, \quad \gamma_2 \mapsto \sigma_2^2.$$

Hence, after performing a Kummer cover of order 3, one obtains:

$$\begin{aligned} \gamma_1 &\mapsto \sigma_1^2 \sigma_3^2 \sigma_5^2 \\ \gamma_1^{\gamma_2} &\mapsto (\sigma_1^2 \sigma_3^2 \sigma_5^2)^{\sigma_4 \sigma_3 \sigma_2} \\ \gamma_1^{\gamma_2^2} &\mapsto (\sigma_1^2 \sigma_3^2 \sigma_5^2)^{\sigma_4^2 \sigma_3 \sigma_2^2} \\ \gamma_2^3 &\mapsto \sigma_4 * (\sigma_3 \sigma_2)^3 \end{aligned}$$

(see (6.2)).

In order to make this monodromy generic at infinity one applies Proposition 4.2 obtaining the new braid:

$$(\Delta_3^2 c^{-1})^3 = ((\sigma_3 \sigma_2)^3)^{\sigma_4^{-1} \sigma_1 \sigma_3 \sigma_5}.$$

Using Proposition 4.9 one can obtain a generic braid monodromy as follows:

$$\begin{aligned} &(\Delta_{5,7}^2, (\Delta_{3,5}^2)^{\Delta_{5,7}^{-1}}, (\Delta_{1,3}^2)^{\Delta_{3,5}^{-1}\Delta_{5,7}^{-1}}, (\Delta_{5,7}^2)^{\sigma_4\sigma_3\sigma_2\sigma_7}, (\Delta_{3,5}^2)^{\Delta_{5,7}^{-1}\sigma_4\sigma_3\sigma_2\sigma_7}, \\ &(\Delta_{1,3}^2)^{\Delta_{3,5}^{-1}\Delta_{5,7}^{-1}\sigma_4\sigma_3\sigma_2\sigma_7}, (\Delta_{5,7}^2)^{\sigma_4^2\sigma_3\sigma_2^2\sigma_7\sigma_8}, (\Delta_{3,5}^2)^{\Delta_{5,7}^{-1}\sigma_4^2\sigma_3\sigma_2^2\sigma_7\sigma_8}, \\ &(\Delta_{1,3}^2)^{\Delta_{3,5}^{-1}\Delta_{5,7}^{-1}\sigma_4^2\sigma_3\sigma_2^2\sigma_7\sigma_8}, \sigma_4 * (\sigma_3\sigma_2)^3, ((\sigma_3\sigma_2)^3)^{\sigma_4^{-1}\sigma_1\sigma_3\sigma_5}). \end{aligned}$$

Another direct application is the monodromy of the MacLane arrangement [29], which is obtained from $\bar{\mathcal{C}}_3$ by deleting one line. This results in deleting one of the strings, that is,

$$\begin{aligned} &(\Delta_{5,7}^2, (\Delta_{3,5}^2)^{\Delta_{5,7}^{-1}}, (\Delta_{1,3}^2)^{\Delta_{3,5}^{-1}\Delta_{5,7}^{-1}}, (\Delta_{5,7}^2)^{\sigma_4\sigma_3\sigma_2\sigma_7}, (\Delta_{3,5}^2)^{\Delta_{5,7}^{-1}\sigma_4\sigma_3\sigma_2\sigma_7}, (\Delta_{1,3}^2)^{\Delta_{3,5}^{-1}\Delta_{5,7}^{-1}\sigma_4\sigma_3\sigma_2\sigma_7}, \\ &(\sigma_5^2)^{\sigma_4^2\sigma_3\sigma_2^2}, (\sigma_3^2)^{\sigma_4^2\sigma_3\sigma_2^2}, (\sigma_1^2)^{\sigma_4^2\sigma_3\sigma_2^2}, \sigma_4 * (\sigma_3\sigma_2)^3, ((\sigma_3\sigma_2)^3)^{\sigma_4^{-1}\sigma_1\sigma_3\sigma_5}). \end{aligned}$$

The computational difficulty of the braid monodromy of the 9-Ceva as well as the MacLane arrangements comes from the fact that they cannot be given by real equations; MacLane arrangement is the smallest one with this property. Using our construction everything is reduced to computing the braid monodromy of a very simple line arrangement. Braid monodromy for generalized Ceva arrangements [10] ($n > 3$) can also be obtained.

6.5. A useful nodal cubic.

Let us consider the following cubic $\bar{\mathcal{C}} = \{f(x, y, z) = (x + y + z)^3 - 27xyz = 0\}$, whose real picture appears in Figure 15(a). The usual projection from P_y has four non-generic fibers \bar{L}_t at $t = \infty, 1, 0$, where \bar{L}_∞ and \bar{L}_0 are tangent lines at inflection points and \bar{L}_1 is the vertical line through the node (which is real, but whose branches are not). For this reason the real picture is not enough to recover the braid monodromy of $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$. However, the extra information required can be obtained from drawing the real part of because of the missing complex conjugate branches (shown in Figure 15(b) as dotted lines).




FIGURE 15.

In order to find these dotted lines one can proceed as follows. Let $p(y) := y^3 - a_1y + a_2y^2 - a_3 \in \mathbb{R}[y]$ and assume it has only one real root, say t_1 . Let $t_2, t_3 \in \mathbb{C}$ be the remaining roots, $t_3 = \bar{t}_2$. Note that their common real part is $\frac{t_2+t_3}{2}$.

Lemma 6.5. *Under the above conditions, the polynomial*

$$q(y) = y^3 - a_1y^2 + \frac{a_1^2 + a_2}{4}y - \frac{a_1a_2 - a_3}{8}$$

contains exactly one real root, which is the common real part $\frac{t_2+t_3}{2}$ of the complex conjugate roots t_2, t_3 of $p(y)$.

Proof. Since $a_i = s_i(t_1, t_2, t_3)$ the symmetric polynomial of degree i , it is enough to show that $a_1 = s_1(\tilde{t}_1, \tilde{t}_2, \tilde{t}_3)$, $\frac{a_1^2 + a_2}{4} = s_2(\tilde{t}_1, \tilde{t}_2, \tilde{t}_3)$, and $\frac{a_1a_2 - a_3}{8} = s_3(\tilde{t}_1, \tilde{t}_2, \tilde{t}_3)$, where $\tilde{t}_i = \frac{t_j + t_k}{2}$, $\{i, j, k\} = \{1, 2, 3\}$, which is a simple exercise. \square

Since the affine cubic $f(x_0, y, 1)$ satisfies the conditions of Lemma 6.5 for any fixed $\bar{x} \in \mathbb{R}$ the real parts of the complex roots of $f(x_0, y, 1)$ are given by the equation:

$$0 = y^3 + 3(x_0+1)y^2 + 3\frac{4x_0^2 - x_0 + 4}{4} + (x_0+1)\frac{8x_0^2 - 65x_0 + 8}{8} = (y+x_0+1)^3 - \frac{27}{4}x_0y - \frac{81}{8}x_0(x_0+1).$$

This is enough to compute the required braid monodromy, see [6] for details.

In order to show the extended braid monodromy $\tilde{\nabla}$ of \mathcal{C} note that L_Y is also tangent to \mathcal{C} at the inflection point $(-1, 0)$. Hence $\tilde{B} = \{1, 0, -1\}$. Choose a geometric basis as in Figure 15(b) and a system diagram on the fiber given by decreasing lexicographic order on \mathbb{C} . For instance, according to Figure 15(b) σ_1 is the half-twist exchanging the two complex conjugated roots of $f(t_0, y, 1)$, σ_2 is the half-twist exchanging the complex conjugated root of $f(t_0, y, 1)$ of negative real part and 0. Then, $\tilde{\nabla} : \pi_1(\mathbb{C} \setminus \tilde{B}; t_0) \rightarrow \mathbb{B}_{3,1}$ is given by:

$$\begin{aligned} \hat{\gamma}_1 &\mapsto \sigma_1^2 \\ \hat{\gamma}_2 &\mapsto (\sigma_2\sigma_3)\sigma_1^{-1}\sigma_2 \\ \hat{\gamma}_3 &\mapsto \sigma_2^6. \end{aligned}$$

In order to meet the hypothesis of §2 one needs the system diagram to be such that the last string is the one that remains constant. This means one needs to conjugate $\tilde{\nabla}$ by σ_3^{-1} . Combining this with a conjugation by σ_2^{-1} , the Hurwitz move h_2^{-1} (changing $\hat{\gamma}_i$ to γ_i), and another conjugation by σ_1^{-1} one obtains:

$$(6.9) \quad \begin{aligned} \gamma_1 &\mapsto \sigma_2^2 \\ \gamma_2 &\mapsto \sigma_2 * \sigma_3^6 \\ \gamma_3 &\mapsto (\sigma_1\sigma_2)\sigma_3^2. \end{aligned}$$

An easy computation gives that the product of Δ^2 and the inverse of the pseudo-Coxeter element is $(\sigma_1\sigma_2)\sigma_3^2\sigma_2$.

6.6. The dual of the nodal quartic.

The dual curve of a nodal quartic is a sextic curve with six cusps and four nodes. Moreover, this curve appears as a generic plane section of the discriminant of the polynomials of degree 4 and its fundamental group has been computed by Zariski [36]: it is the braid group on \mathbb{S}^2 with four strings.

This sextic can also be obtained from the cubic in §6.5 using a Kummer cover π_2 . Each inflection point in the axes produces two cusps while the double point produces four nodes, see Example 5.5.

In order to give a braid monodromy for $(\mathcal{C}_2, P_y, \bar{L}_\infty)$ from (6.9) one needs to combine $\hat{\rho}'_2$ (see Example 3.8) and Lemma 2.7. We start with the first part of (2.2):

$$(6.10) \quad \begin{array}{ccccccc} \mathbb{F}_5 & \hookrightarrow & \mathbb{F}_3 & \xrightarrow{\bar{\nabla}} & \mathbb{B}_{3,1} \\ \tilde{\gamma}_1 & \mapsto & \gamma_1 & \mapsto & \sigma_2^2 \\ \tilde{\gamma}_2 & \mapsto & \gamma_2 & \mapsto & \sigma_2 * \sigma_3^6 \\ \tilde{\gamma}_3 & \mapsto & \gamma_1^3 & \mapsto & (\sigma_2^2)^{\sigma_3^{-2} \sigma_1 \sigma_2 \sigma_3^2} = (\sigma_2 \sigma_3^2 \sigma_1) * \sigma_2^2 \\ \tilde{\gamma}_4 & \mapsto & \gamma_2^3 & \mapsto & (\sigma_3^6)^{\sigma_2^{-1} \sigma_3^{-2} \sigma_1 \sigma_2 \sigma_3^2} = (\sigma_3^6)^{\sigma_2^{-1} \sigma_1} \\ \tilde{\gamma}_5 & \mapsto & \gamma_3^2 & \mapsto & \sigma_3^{-2} * (\sigma_1 \sigma_2)^2 \end{array}$$

Using $\hat{\rho}'_2$ we obtain the braid monodromy of $(\bar{\mathcal{C}}_2, P_y, \bar{L}_\infty)$:

$$(6.11) \quad (\hat{\sigma}_2^2 \hat{\sigma}_4^2, (\hat{\sigma}_2 \hat{\sigma}_4) * \hat{\sigma}_3^3, (\hat{\sigma}_2 \hat{\sigma}_4 \hat{\sigma}_3 \hat{\sigma}_5 \hat{\sigma}_1) * (\hat{\sigma}_2^2 \hat{\sigma}_4^2), (\hat{\sigma}_3^3)^{\hat{\sigma}_2^{-1} \hat{\sigma}_4^{-1} \hat{\sigma}_5 \hat{\sigma}_1}, ((\hat{\sigma}_1 \hat{\sigma}_2)^2 (\hat{\sigma}_5 \hat{\sigma}_4)^2)^{\hat{\sigma}_3}).$$

Using Proposition 4.2, the monodromy can be made generic at infinity adding the image by $\hat{\rho}'_2$ of $(\sigma_2^{-1} \sigma_3^2) * (\sigma_2 \sigma_1)^3$ which is $((\hat{\sigma}_1 \hat{\sigma}_2)^2 (\hat{\sigma}_5 \hat{\sigma}_4)^2)^{\hat{\sigma}_3 \hat{\sigma}_2 \hat{\sigma}_4}$.

In order to obtain a generic braid monodromy we apply Propositions 4.3 and 4.4. Table 1 shows the decompositions:.

$\hat{\sigma}_2^2 \hat{\sigma}_4^2$	(σ_2^2, σ_4^2)
$(\hat{\sigma}_2 \hat{\sigma}_4) * \hat{\sigma}_3^3$	$((\hat{\sigma}_2 \hat{\sigma}_4) * \hat{\sigma}_3^3)$
$(\hat{\sigma}_2 \hat{\sigma}_4 \hat{\sigma}_3 \hat{\sigma}_5 \hat{\sigma}_1) * (\hat{\sigma}_2^2 \hat{\sigma}_4^2)$	$((\hat{\sigma}_2 \hat{\sigma}_4 \hat{\sigma}_3 \hat{\sigma}_1) * (\hat{\sigma}_2^2), (\hat{\sigma}_2 \hat{\sigma}_4 \hat{\sigma}_3 \hat{\sigma}_5) * (\hat{\sigma}_4^2))$
$(\hat{\sigma}_3^3)^{\hat{\sigma}_2^{-1} \hat{\sigma}_4^{-1} \hat{\sigma}_5 \hat{\sigma}_1}$	$((\hat{\sigma}_3^3)^{\hat{\sigma}_2^{-1} \hat{\sigma}_4^{-1} \hat{\sigma}_5 \hat{\sigma}_1})$
$((\hat{\sigma}_1 \hat{\sigma}_2)^2 (\hat{\sigma}_5 \hat{\sigma}_4)^2)^{\hat{\sigma}_3}$	$(\hat{\sigma}_2^{\hat{\sigma}_1 \hat{\sigma}_3}, \hat{\sigma}_1^3, \hat{\sigma}_4^{\hat{\sigma}_5 \hat{\sigma}_3}, \hat{\sigma}_5^3)$
$((\hat{\sigma}_1 \hat{\sigma}_2)^2 (\hat{\sigma}_5 \hat{\sigma}_4)^2)^{\hat{\sigma}_3 \hat{\sigma}_2 \hat{\sigma}_4}$	$(\hat{\sigma}_2^{\hat{\sigma}_1 \hat{\sigma}_3 \hat{\sigma}_2 \hat{\sigma}_4}, (\hat{\sigma}_1^3)^{\hat{\sigma}_2}, \hat{\sigma}_4^{\hat{\sigma}_5 \hat{\sigma}_3 \hat{\sigma}_2 \hat{\sigma}_4}, (\hat{\sigma}_5^3)^{\hat{\sigma}_4})$

TABLE 1. From generic at infinity to generic braid monodromy

We can compute the fundamental group of $\mathbb{C}^2 \setminus \mathcal{C}_2$. It is well known [36] that this group is obtained as a central extension of the braid group of \mathbb{S}^2 with four strings by \mathbb{Z} . More precisely, applying Theorem 1.7 (without the relation $\mu_6 \cdots \mu_1 = 1$) we obtain a group with generators μ_1, \dots, μ_6 and relators

$$\begin{aligned} [\mu_4, \mu_5] &= [\mu_2^2, \mu_5 \mu_4^{-1}] = 1, \mu_2 \mu_4 \mu_2 = \mu_4 \mu_2 \mu_4, \mu_2 \mu_5 \mu_2 = \mu_5 \mu_2 \mu_5, \\ \mu_6 &= \mu_2^{\mu_5}, \mu_1 = \mu_4 * \mu_2, \mu_3 = \mu_5^{\mu_2 \mu_4}, \underbrace{1 = 1}_{7 \text{ times}}. \end{aligned}$$

This presentation is obtained using **GAP4** [22].

From the original presentation we obtain this one only using Tietze moves of type I and II. Hence, by Libgober's result [27], one can verify that $\mathbb{C}^2 \setminus \mathcal{C}_2$ has the homotopy type of $K \vee \bigvee_{j=1}^7 \mathbb{S}^2$, where K is the 2-complex associated with the presentation

$$\langle x, y, z \mid [x, z] = [y^2, xz^{-1}] = 1, xyx = yxy, yzy = zyz \rangle.$$

6.7. Hesse arrangement.

It can be obtained from the cubic in §6.5 using a Kummer cover π_3 : it is the preimage of the cubic and the three ramification lines. It can be seen using Proposition 4.3 and Example 5.5, and also directly from the equations:

$$(x^3 + y^3 + z^3)^3 - 27x^3y^3z^3 = \prod_{\zeta_1^3=1} (x^3 + y^3 + z^3 - 3\zeta_1xyz).$$

Hence $xyz = 0$ and the above factors give the four reducible members of the pencil of cubics generated by $x^3 + y^3 + z^3 = 0$ and $xyz = 0$. The base points of this pencil are the nine common inflection points of the irreducible cubics of the pencil. The reducible fibers split as product of lines:

$$x^3 + y^3 + z^3 - 3\zeta_1xyz = \prod_{\zeta_2^3=1} (x + \zeta_2y + \zeta_1\bar{\zeta}_2z).$$

These are the lines joining the inflection points, i.e., we obtain Hesse arrangement. We start with the braid monodromy with respect to $(\bar{\mathcal{C}}_3, P_y, \bar{L}_\infty)$: $(\sigma_2^2, \sigma_2 * \sigma_3^6, (\sigma_1\sigma_2)^{\sigma_3^2})$, and the braid at infinity is $(\sigma_1\sigma_2)^{\sigma_3^2\sigma_2}$. In order to simplify the braid monodromy of the Hesse arrangement we conjugate by $\sigma_3^{-2}\sigma_2^{-1}$ and we obtain:

$$((\sigma_2^2)^{\sigma_3^2}, \sigma_3^6, \sigma_2\sigma_1), \quad \infty \mapsto (\sigma_2\sigma_3^2\sigma_2^{-1}\sigma_3^{-2}) * (\sigma_1\sigma_2).$$

We proceed as in §6.6 but we perform a Hurwitz move to the base of Lemma 2.7 in order to obtain simpler braids:

$$(6.12) \quad \begin{array}{ccccccc} \mathbb{F}_7 & \hookrightarrow & \mathbb{F}_3 & \xrightarrow{\tilde{\nabla}} & \mathbb{B}_{3,1} \\ \tilde{\gamma}_1 & \mapsto & \gamma_1 & \mapsto & (\sigma_2^2)^{\sigma_3^2} \\ \tilde{\gamma}_2 & \mapsto & \gamma_2 & \mapsto & \sigma_3^6 \\ \tilde{\gamma}_3 & \mapsto & \gamma_1^{\gamma_3} & \mapsto & (\sigma_2^2)^{\sigma_3^2\sigma_2\sigma_1} \\ \tilde{\gamma}_4 & \mapsto & \gamma_2^{\gamma_3} & \mapsto & (\sigma_3^6)^{\sigma_2\sigma_1} \\ \tilde{\gamma}_5 & \mapsto & \gamma_3^3 & \mapsto & (\sigma_2\sigma_1)^3 \\ \tilde{\gamma}_6 & \mapsto & \gamma_3 * \gamma_1 & \mapsto & (\sigma_2\sigma_1\sigma_3^{-2}) * \sigma_2^2 \\ \tilde{\gamma}_7 & \mapsto & \gamma_3 * \gamma_2 & \mapsto & \sigma_2 * \sigma_3^6. \end{array}$$

Let us recall the map $\tilde{\rho}_3 : \mathbb{B}_{3,1} \rightarrow \mathbb{B}_{9,1}$ of Proposition 3.7:

$$(6.13) \quad \begin{array}{ll} \sigma_1 & \mapsto \tilde{\sigma}_{1,1}\tilde{\sigma}_{1,2}\tilde{\sigma}_{1,3} \\ \sigma_2 & \mapsto \tilde{\sigma}_{2,1}\tilde{\sigma}_{2,2}\tilde{\sigma}_{2,3} \\ \sigma_3^2 & \mapsto \tilde{\sigma}_{3,3}^2\tilde{\sigma}_{2,3}\tilde{\sigma}_{1,3}\tilde{\sigma}_{3,2}\tilde{\sigma}_{2,2}\tilde{\sigma}_{1,2}\tilde{\sigma}_{3,1}\tilde{\sigma}_{1,2}^{-1}\tilde{\sigma}_{2,2}^{-1}\tilde{\sigma}_{1,3}^{-1}\tilde{\sigma}_{2,3}^{-1}. \end{array}$$

For convenience we rewrite this map in the usual generators:

$$(6.14) \quad \begin{aligned} \sigma_1 &\mapsto \sigma_1 \sigma_4 \sigma_7 \\ \sigma_2 &\mapsto \sigma_2 \sigma_5 \sigma_8 \\ \sigma_3^2 &\mapsto \sigma_9^2 \sigma_8 \sigma_7 \sigma_6 \sigma_5 \sigma_4 \sigma_3 \sigma_4^{-1} \sigma_5^{-1} \sigma_7^{-1} \sigma_8^{-1} = (\sigma_9^2 \sigma_8 \sigma_7)^{\sigma_6 \sigma_5 \sigma_4 \sigma_3 \sigma_7 \sigma_6}. \end{aligned}$$

It is worthwhile to note that $(\sigma_9^2 \sigma_8 \sigma_7)^3 = \Delta_{7,10}^2$. Let us denote $\tau := \sigma_6 \sigma_5 \sigma_4 \sigma_3 \sigma_7 \sigma_6$ and $\Theta := (\sigma_9^2 \sigma_8 \sigma_7)^\tau$. Next step is to write down the braid monodromy of $(\bar{\mathcal{C}}_3, P_y, \bar{L}_\infty)$:

$$\begin{aligned} &((\sigma_2^2 \sigma_5^2 \sigma_8^2)^\Theta, (\Delta_{7,10}^2)^\tau, (\sigma_2^2 \sigma_5^2 \sigma_8^2)^{\Theta \sigma_2 \sigma_1 \sigma_5 \sigma_4 \sigma_8 \sigma_7}, (\Delta_{7,10}^2)^{\tau \sigma_2 \sigma_1 \sigma_5 \sigma_4 \sigma_8 \sigma_7}, \Delta_{1,3}^2 \Delta_{4,6}^2 \Delta_{7,9}^2, \\ &(\sigma_2 \sigma_1 \sigma_5 \sigma_4 \sigma_8 \sigma_7 \Theta^{-1}) * (\sigma_2^2 \sigma_5^2 \sigma_8^2), (\sigma_2 \sigma_5 \sigma_8 \tau^{-1}) * \Delta_{7,10}^2). \end{aligned}$$

This monodromy becomes generic at infinity adding $(\sigma_2 \sigma_5 \sigma_8 \Theta \sigma_8^{-1} \sigma_5^{-1} \sigma_2^{-1} \Theta^{-1}) * (\Delta_{1,3}^2 \Delta_{4,6}^2 \Delta_{7,9}^2)$. Let us denote $\eta := \sigma_2 \sigma_5 \sigma_8 \Theta \sigma_8^{-1} \sigma_5^{-1} \sigma_2^{-1} \Theta^{-1}$.

To compute a generic braid monodromy to this arrangement one has to apply Proposition 4.9 where we have two vertical lines corresponding to the fifth and eighth braids:

$$(6.15) \quad \begin{aligned} &((\sigma_2^2)^\Theta, (\sigma_5^2)^\Theta, (\sigma_8^2)^\Theta, (\Delta_{7,10}^2)^\tau, (\sigma_2^2)^{\Theta \sigma_2 \sigma_1 \sigma_5 \sigma_4 \sigma_8 \sigma_7}, (\sigma_5^2)^{\Theta \sigma_2 \sigma_1 \sigma_5 \sigma_4 \sigma_8 \sigma_7}, (\sigma_8^2)^{\Theta \sigma_2 \sigma_1 \sigma_5 \sigma_4 \sigma_8 \sigma_7}, \\ &(\Delta_{7,10}^2)^{\tau \sigma_2 \sigma_1 \sigma_5 \sigma_4 \sigma_8 \sigma_7}, \sigma_{10}^2, \sigma_{10} * \Delta_{7,10}^2, (\sigma_{10} \Delta_{7,10}) * \Delta_{4,7}^2, (\sigma_{10} \Delta_{7,10} \Delta_{4,7}) * \Delta_{1,4}^2, \\ &(\sigma_2 \sigma_1 \sigma_5 \sigma_4 \sigma_8 \sigma_7 \Theta^{-1}) * \sigma_2^2, (\sigma_2 \sigma_1 \sigma_5 \sigma_4 \sigma_8 \sigma_7 \Theta^{-1}) * \sigma_5^2, (\sigma_2 \sigma_1 \sigma_5 \sigma_4 \sigma_8 \sigma_7 \Theta^{-1}) * \sigma_8^2, \\ &(\sigma_2 \sigma_5 \sigma_8 \tau^{-1}) * \Delta_{7,10}^2, (\sigma_{11}^{-1} \eta) * \sigma_{10}^2, (\sigma_{11}^{-1} \eta \sigma_{10}) * \Delta_{7,10}^2, (\sigma_{11}^{-1} \eta \sigma_{10} \Delta_{7,10}) * \Delta_{4,7}^2, \\ &(\sigma_{11}^{-1} \eta \sigma_{10} \Delta_{7,10} \Delta_{4,7}) * \Delta_{1,4}^2, \sigma_{11}^2). \end{aligned}$$

The braid monodromy was also obtained (computer-aid) in [17].

6.8. A smooth cubic and its inflection points and sextics with six cusps outside a conic.

The Kummer cover π_3 of the curve $\bar{\mathcal{C}}$ of equation $(x+y+z)(x+y)(y+z)(x+z) = 0$ produces a Fermat cubic (as the preimage of the line $x+y+z=0$) while the other three lines become the tangent lines to the nine inflection points. The extended braid monodromy of $(\bar{\mathcal{C}}, P_y, \bar{L}_\infty)$ is




FIGURE 16. Arrangement $\bar{\mathcal{C}}$.

given by $(\sigma_2^2, \sigma_2 * (\sigma_1^2 \sigma_3^2), (\sigma_2 \sigma_3 \sigma_2^{-1}) * \sigma_1^2)$. In order to meet the hypothesis of §2 one needs both that the last generator in the ordered geometric basis corresponds to $x = 0$ and that the system diagram is such that the last string is the one that remains constant. One can meet the first requirement by using the Hurwitz move h_2^{-1} :

$$(\sigma_2^2, (\sigma_2 \sigma_3^{-1} \sigma_1^{-1}) * \sigma_2^2, \sigma_2 * (\sigma_1^2 \sigma_3^2)).$$

In order to switch the string $y = 0$ to the last position one must perform the automorphism $\sigma_i \mapsto \sigma_{4-i}$, but these braids are invariant by the morphism. After these changes the braid around infinity is $(\sigma_1^2 \sigma_3^2)^{\sigma_2}$.

Using Proposition 3.7 one can apply (6.13). In [3] sextics with six cusps not on a line have been constructed by means of 3rd Kummer cover where the three lines are tangent lines to a smooth cubic at non-aligned inflection points. Hence we will be interested only in the three tangent lines with these property, namely, two preimages of $x + z = 0$, one preimage of $y + z = 0$ and none of $x + y = 0$. Hence, we can forget the second string (recall the automorphism) and we obtain (the first braid disappears since it is trivial):

$$((\sigma_2^2)^{\sigma_1}, \sigma_1^2)$$

and hence the braid around infinity is σ_2^2 . We have to forget some strings in the mapping $\hat{\rho}_3$. This causes the morphism to not be well defined in $\mathbb{B}_{2,1}$, but only in the pure braid group:

$$\begin{array}{ccc} \sigma_1^2 & \mapsto & \sigma_1^2 \\ \sigma_2^2 & \mapsto & 1 \\ (\sigma_2^2)^{\sigma_1} & \mapsto & (\sigma_3 \sigma_2)^{\sigma_1}. \end{array}$$

Hence the braid monodromy of the cubic with the tangent lines is given by:

$$((\sigma_3 \sigma_2)^{\sigma_1}, (\sigma_3 \sigma_2)^{\sigma_1^3}, (\sigma_3 \sigma_2)^{\sigma_1^5}, \sigma_1^6);$$

where the first two braids correspond to the tangent lines in the curve while the second string corresponds to the remaining tangent line. Some operations are need: conjugation by σ_1^{-3} , cyclic permutation, permutation of the order of the strings and forgetting one braid (at infinity):

$$(\sigma_3^6, \sigma_3^2 * (\sigma_1 \sigma_2), \sigma_1 \sigma_2),$$

the braid at infinity being $(\sigma_1 \sigma_2)^{\sigma_3^2}$. This result coincides with the one obtained using Carmona's program [12].

We first consider the braid monodromy factorization induced by this braid monodromy after a double cover of the base, that is, $\mathbb{F}_5 \hookrightarrow \mathbb{F}_3 \rightarrow \mathbb{B}_{3,1}$:

$$(\sigma_3^6, \sigma_3^2 * (\sigma_1 \sigma_2), (\sigma_3^6)^{\sigma_2}, (\sigma_2^{-1} \sigma_3^2) * (\sigma_1 \sigma_2), (\sigma_1 \sigma_2)^2, ((\sigma_1 \sigma_2)^2)^{\sigma_3^2}).$$

Then perform the morphism $\tilde{\rho}$ (see Example 3.8) and a generic deformation, resulting in:

$$(\sigma_3^3, \sigma_1, \sigma_5, \sigma_3^{\sigma_2}, \sigma_3^{\sigma_4}, (\sigma_3^3)^{\sigma_2 \sigma_4}, \sigma_1^{\sigma_2}, \sigma_5^{\sigma_4}, \sigma_3^{\sigma_2^2 \sigma_4}, \sigma_3^{\sigma_2 \sigma_4^2}, \sigma_2^{\sigma_1}, \sigma_1^3, \sigma_4^{\sigma_5}, \sigma_5^3, \sigma_2^{\sigma_1 \sigma_3}, \sigma_1^3, \sigma_4^{\sigma_5 \sigma_3}, \sigma_5^3).$$

A straightforward computation shows that $\pi_1(\mathbb{C}^2 \setminus \mathcal{C}_2) = \mathbb{Z}$. Using [27] one can check that

$$\mathbb{C}^2 \setminus \mathcal{C}_2 \simeq \mathbb{S}^1 \vee \bigvee_{i=1}^{13} \mathbb{S}^2.$$

REFERENCES

- [1] M. Amram, C. Ciliberto, R. Miranda, and M. Teicher, *Braid monodromy factorization for a non-prime K3 surface branch curve*, Israel J. Math. **170** (2009), 61–93.
- [2] M. Amram and M. Teicher, *On the degeneration, regeneration and braid monodromy of $T \times T$* , Acta Appl. Math. **75** (2003), no. 1-3, 195–270, Monodromy and differential equations (Moscow, 2001).
- [3] E. Artal, *Sur les couples de Zariski*, J. Algebraic Geom. **3** (1994), no. 2, 223–247.
- [4] E. Artal and J. Carmona, *Zariski pairs, fundamental groups and Alexander polynomials*, J. Math. Soc. Japan **50** (1998), no. 3, 521–543.
- [5] E. Artal, J. Carmona, and J.I. Cogolludo-Agustín, *Braid monodromy and topology of plane curves*, Duke Math. J. **118** (2003), no. 2, 261–278.
- [6] E. Artal, J. Carmona, J.I. Cogolludo-Agustín, and H. Tokunaga, *Sextics with singular points in special position*, J. Knot Theory Ramifications **10** (2001), no. 4, 547–578.
- [7] E. Artal, J.I. Cogolludo-Agustín, and H. Tokunaga, *A survey on Zariski pairs*, Algebraic geometry in East Asia—Hanoi 2005, Adv. Stud. Pure Math., vol. 50, Math. Soc. Japan, Tokyo, 2008, pp. 1–100.
- [8] W.A. Arvola, *The fundamental group of the complement of an arrangement of complex hyperplanes*, Topology **31** (1992), no. 4, 757–765.
- [9] D. Auroux and L. Katzarkov, *Branched coverings of \mathbb{CP}^2 and invariants of symplectic 4-manifolds*, Invent. Math. **142** (2000), no. 3, 631–673.
- [10] G. Barthel, F. Hirzebruch, and T. Höfer, *Geradenkonfigurationen und Algebraische Flächen*, Aspects of Mathematics, D4, Friedr. Vieweg & Sohn, Braunschweig, 1987.
- [11] D. Bessis and J. Michel, *Explicit presentations for exceptional braid groups*, Experiment. Math. **13** (2004), no. 3, 257–266.
- [12] J. Carmona, *Monodromía de trenzas de curvas algebraicas planas*, Ph.D. thesis, Universidad de Zaragoza, 2003.
- [13] F. Catanese, M. Lönne, and B. Wajnryb, *Moduli spaces of surfaces and monodromy invariants*, Proceedings of the Gökova Geometry-Topology Conference 2009, Int. Press, Somerville, MA, 2010, pp. 58–98.
- [14] F. Catanese and B. Wajnryb, *The 3-cuspidal quartic and braid monodromy of degree 4 coverings*, Projective varieties with unexpected properties, Walter de Gruyter GmbH & Co. KG, Berlin, 2005, pp. 113–129.
- [15] O. Chisini, *Una suggestiva rappresentazione reale per le curve algebriche piane*, Ist. Lombardo, Rend., II. Ser. **66** (1933), 1141–1155.
- [16] J.I. Cogolludo-Agustín, *Fundamental group for some cuspidal curves*, Bull. London Math. Soc. **31** (1999), no. 2, 136–142.
- [17] ———, *Braid monodromy of algebraic curves*, Ann. Math. Blaise Pascal **18** (2011), no. 1, 141–209.
- [18] J.I. Cogolludo-Agustín and R. Kloosterman, *Mordell-Weil groups and Zariski triples*, Geometry and Arithmetic (G. Farkas C. Faber and R. de Jong, eds.), EMS Congress Reports, Europ. Math. Soc., 2012, Also available at [arXiv:1111.5703](https://arxiv.org/abs/1111.5703) [math.AG].
- [19] D.C. Cohen and A.I. Suciu, *The braid monodromy of plane algebraic curves and hyperplane arrangements*, Comment. Math. Helv. **72** (1997), no. 2, 285–315.
- [20] R. Cordovil and J. L. Fachada, *Braid monodromy groups of wiring diagrams*, Boll. Un. Mat. Ital. B (7) **9** (1995), no. 2, 399–416.
- [21] I. Dolgachev and A. Libgober, *On the fundamental group of the complement to a discriminant variety*, Algebraic geometry (Chicago, Ill., 1980), Lecture Notes in Math., vol. 862, Springer, Berlin, 1981, pp. 1–25.

- [22] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.4*, 2004, available at (<http://www.gap-system.org>).
- [23] A. Hirano, *Construction of plane curves with cusps*, Saitama Math. J. **10** (1992), 21–24.
- [24] E.R. van Kampen, *On the fundamental group of an algebraic curve*, Amer. J. Math. **55** (1933), 255–260.
- [25] V.S. Kulikov, *Generic coverings of the plane and braid monodromy invariants*, The Fano Conference, Univ. Torino, Turin, 2004, pp. 533–558.
- [26] V.S. Kulikov and M. Teicher, *Braid monodromy factorizations and diffeomorphism types*, Izv. Ross. Akad. Nauk Ser. Mat. **64** (2000), no. 2, 89–120.
- [27] A. Libgober, *On the homotopy type of the complement to plane algebraic curves*, J. Reine Angew. Math. **367** (1986), 103–114.
- [28] M. Lönne, *Fundamental groups of projective discriminant complements*, Duke Math. J. **150** (2009), no. 2, 357–405.
- [29] S. MacLane, *Some Interpretations of Abstract Linear Dependence in Terms of Projective Geometry*, Amer. J. Math. **58** (1936), no. 1, 236–240.
- [30] B.G. Moishezon, *Stable branch curves and braid monodromies*, Algebraic geometry (Chicago, Ill., 1980), Lecture Notes in Math., vol. 862, Springer, Berlin, 1981, pp. 107–192.
- [31] B.G. Moishezon and M. Teicher, *Braid group techniques in complex geometry. IV. Braid monodromy of the branch curve S_3 of $V_3 \rightarrow \mathbb{CP}^2$ and application to $\pi_1(\mathbb{CP}^2 - S_3, *)$* , Classification of algebraic varieties (L'Aquila, 1992), Contemp. Math., vol. 162, Amer. Math. Soc., Providence, RI, 1994, pp. 333–358.
- [32] L. Rudolph, *Algebraic functions and closed braids*, Topology **22** (1983), no. 2, 191–202.
- [33] M. Salvetti, *Arrangements of lines and monodromy of plane curves*, Compositio Math. **68** (1988), no. 1, 103–122.
- [34] A.M. Uludağ, *More Zariski pairs and finite fundamental groups of curve complements*, Manuscripta Math. **106** (2001), no. 3, 271–277.
- [35] O. Zariski, *On the problem of existence of algebraic functions of two variables possessing a given branch curve*, Amer. J. Math. **51** (1929), 305–328.
- [36] ———, *On the Poincaré group of rational plane curves*, Amer. J. Math. **58** (1936), 607–619.
- [37] ———, *The topological discriminant group of a riemann surface of genus p* , Amer. J. Math. **59** (1937), 335–358.

DEPARTAMENTO DE MATEMÁTICAS, IUMA, UNIVERSIDAD DE ZARAGOZA, C. PEDRO CERBUNA 12, 50009 ZARAGOZA, SPAIN

E-mail address: artal@unizar.es, jicogo@unizar.es

CENTRO UNIVERSITARIO DE LA DEFENSA-IUMA, ACADEMIA GENERAL MILITAR, CTRA. DE HUESCA S/N., 50090, ZARAGOZA, SPAIN

E-mail address: jortigas@unizar.es